arXiv:1211.6684v1 [math.AG] 28 Nov 2012

OVERCONVERGENT SUBANALYTIC SUBSETS IN THE FRAMEWORK OF BERKOVICH SPACES

FLORENT MARTIN

Abstract. We study the class of overconvergent subanalytic subsets of a k-affinoid space X when k is a non-archimedean field. These are the images along the projection $X \times \mathbb{B}^n \to X$ of subsets defined with inequalities between functions of $X \times \mathbb{B}^n$ which are overconvergent in the variables of $\mathbb{B}^n$. In particular, we study the local nature, with respect to X, of overconvergent subanalytic subsets. We show that they behave well with respect to the Berkovich topology, but not to the G-topology. This gives counter-examples to previous results on the subject, and a way to correct them. Moreover, we study the case $\dim(X) = 2$, for which a simpler characterisation of overconvergent subanalytic subsets is proven.

Contents

Introduction 1
1. Overconvergent constructible subsets 5
1.1. Constructible data 5
1.2. Overconvergent subanalytic subsets 9
1.3. Weierstrass preparation 10
1.4. Equivalence of the two notions 14
1.5. From a global to a local definition 19
2. Study of various classes 22
2.1. Many families 22
2.2. Rigid semianalytic subsets are not necessarily overconvergent constructible 23
2.3. The other inequalities 25
2.4. Berkovich points versus rigid points 27
3. Overconvergent subanalytic subsets when $\dim(X) = 2$ 28
References 34

Introduction

Notations. In this text k will be a non-trivially valued, non-archimedean field (that is to say a field equipped with a complete non-archimedean non-trivial valuation).
We'll denote by $\mathbb{B}$ (resp. $\mathbb{B}_r$ for a real number r) the closed unit ball of radius 1 (resp. r), seen as a k-analytic space.

Motivations. Since non-archimedean fields are totally disconnected, one can not define the notion of analytic spaces over k as easily as in the case of $\mathbb{R}$ or $\mathbb{C}$. Tate [22] developed such a theory, and called his spaces rigid spaces, whose building blocks are affinoid spaces. However, these spaces are not endowed with a classical topology, but with a Grothendieck topology. Afterwards, V. Berkovich developed another viewpoint for k-analytic geometry [1, 2]. His spaces, called k-analytic spaces or Berkovich spaces, have more points than the corresponding rigid spaces and moreover are equipped with a topology which is locally arcwise connected, moreover, in this theory, affinoid spaces are compact. R. Huber also also developed another view point, in the setting of adic spaces [13], finally there exists also an approach, initiated by Raynaud, using formal geometry (see [4] for instance).
In this text, we'll work in the framework of Berkovich spaces.

Date: November 29, 2012.

Definition 0.1. Let $\mathcal{P}$ be the data, for each affinoid space X, of a family $\mathcal{P}_X$ of subsets of X. If S is a subset of an affinoid space X, we'll say that S has (or satisfies) the property $\mathcal{P}$ if $S \in \mathcal{P}_X$. We'll say that

- the property $\mathcal{P}$ is a G-local property if for all affinoid space X and any subset S of X, S satisfies the property $\mathcal{P}$ if and only if for all finite affinoid covering $\{X_i\}$ of X, $S \cap X_i$ satisfies the property $\mathcal{P}$ relatively to X_i (i.e. $S \cap X_i \in \mathcal{P}_{X_i}$).
- the property $\mathcal{P}$ is a local property if for all affinoid space X and any subset S of X, $S \in \mathcal{P}_X$ if and only if for all $x \in X$, there exists an affinoid neighbourhood U of x such that $S \cap U \in \mathcal{P}_U$.

Note that using the compactness of affinoid spaces, saying that $\mathcal{P}$ is a local property is equivalent to require that for all affinoid space X and any $S \subseteq X$, S satisfies $\mathcal{P}$ if and only if for any finite affinoid covering $\{X_i\}$ of X such that $\{\overset{\circ}{X_i}\}$ is also a covering of X, then $S \cap X_i \in \mathcal{P}_{X_i}$.
As a consequence, if $\mathcal{P}$ is a G-local property, then it is a local property.

Example 0.2. A consequence of Kiehl's theorem [3, 9.4.3] is that the class of Zariski-closed subsets of affinoid spaces defines a class which is G-local.

If X is a k-analytic space, one natural question is to understand which shapes subsets of the form $f(Y)$ have, where $f : Y \to X$ is an analytic map. More precisely, one would like to characterise such images $f(Y)$ (actually rather finite boolean combination of such sets) using only data and functions which are internal to X. Note that this defines a G-local property. Such a question is for instance inspired by Chevalley's theorem [12, II exercise 3.19] which asserts that if $\mathcal{X}$ is a noetherian scheme, images $f(\mathcal{Y})$, where $f : \mathcal{Y} \to \mathcal{X}$ is a morphism of finite type, are precisely the constructible subsets of $\mathcal{X}$ (i.e. in that case, finite union of locally closed subsets of $\mathcal{X}$ for the Zarisky topology).
If we come back to k-analytic spaces, trying to describe subsets $f(Y)$ without assumption on X and Y is impossible. One reasonable restriction is to assume that X is affinoid and restrict to maps $f : Y \to X$ where Y is also affinoid. In this context let us explain the first natural approach:

Definition 0.3. Let X be an affinoid space whose affinoid algebra is $\mathcal{A}$. A subset S of X is called *globally semianalytic* if it is a finite boolean combination of sets of the form $\{x \in X \mid |f(x)| \leq |g(x)|\}$ where f and $g \in \mathcal{A}$ (by finite boolean combination, we mean finitely many use of the set-theoretical operators $\cap$, $\cup$ and ^{c}).

It is easy to check that the class of globally semianalytic subsets properly contains the class of Zariski-closed subsets. It is also also easy to see that it is contained in the class of finite boolean combination of images $f(Y)$, and it is properly contained in it. Indeed, the class of globally semianalytic subsets is not G-local [17, 6.2] (in fact neither local, see Proposition 2.6) whereas the second one is. In fact, the class of finite boolean combination of images $f(Y)$ can alternatively be defined as the class of the projections $\pi(T)$ where $\pi : X \times \mathbb{B}^n \to X$ is the natural projection and T is a globally semianalytic subset of $X \times \mathbb{B}^n$ [18, model completeness and subanalytic sets, Theorem 5.2]. So that we arrive to the following question:
How to describe the subsets of X which are the projections of globally semianalytic subsets of $X \times \mathbb{B}^n$?
In [16,18] Lipshitz and Robinson define and study a bigger class of subsets that they call *subanalytic subsets*, which contains the class we were just discussing. But to our knowledge, it is not known if these two classes coincide. In [10, 11] a result stating an internal description of such images $f(Y)$ as D-semianalytic sets was claimed. But as pointed out by Lipshitz and Robinson [17] the proof contains a mistake, so that the characterisation of projection of globally semianalytic subsets as D-semianalytic subsets remains an open problem. Let us mention that these two approaches share a same idea which is to use the division operator D introduced in [6]. In this paper we'll study an intermediate class. Roughly speaking we'll focus our study on images $f(Y) \subseteq X$ where $f : Y \to X$ is a map of affinoid spaces, with the extra condition that f is overconvergent.

Some results of the paper.

Definition. *Let X be an affinoid space, a subset S of X is an overconvergent subanalytic subset of X if there exist n an integer, T a globally semianalytic subset of $X \times \mathbb{B}^n$ such that $S = \pi(T)$*

where $\pi : X \times \mathbb{B}^n \to X$ *is the natural projection, and such that the functions used to define* T *are overconvergent with respect to the variables of* $\mathbb{B}^n$.

This class of subsets has been initially given and studied in [19, 1.3.8]. At first sight, this definition is not local, so one can naturally wonder if this a local or G-local property, in the sense of definition 0.1. We give the following answers:

Theorem. *2.4 Being an overconvergent subanalytic subset is not a* G*-local property.*

This contradicts some of the results of [19]. But this can been corrected with the use of Berkovich topology:

Theorem. *1.35 Being an overconvergent subanalytic subset is a local property.*

Thanks to this local property, we can unambiguously give the following definition:

Definition. *Let* X *be a good* k*-analytic space. A subset* S *of* X *is overconvergent subanalytic if for all* $x \in X$ *there exists an affinoid neighbourhood* V *of* x *such that* $S \cap V$ *is overconvergent subanalytic in* V *(according to the definition we have given above).*

In the same way:

Definition. *Let* X *be a good* k*-analytic space. A subset* S *of* X *is overconvergent subanalytic if for all* $x \in X$ *there exists an affinoid neighbourhood* V *of* x *such that* $S \cap V$ *is overconvergent subanalytic in* V *(according to the definition we have given above).*

As we've mentioned above, the class of globally semianalytic subset is not local (proposition 2.6). We then give the following definition:

Definition. *Let* X *be a good* k*-analytic space. A subset* S *of* X *is locally semianalytic if for all* $x \in X$ *there exists an affinoid neighbourhood* V *of* x *such that* $S \cap V$ *is globally semianalytic in* V *.*

Theorem. *3.13 Let* k *be algebraically closed, and* X *be a quasi-smooth (equivalently rig-smooth, or regular)* k*-analytic space of dimension 2. Then a subset* S *of* X *is overconvergent subanalytic if and only if it is locally semianalytic.*

Let us briefly explain what will be the main ingredient of the proofs. We'll introduce a new class of subsets of X that we'll call overconvergent constructible subsets of X. Then we'll show that:

Theorem. *1.31* OVERCONVERGENT CONSTRUCTIBLE = OVERCONVERGENT SUBANALYTIC.

Overconvergent constructible subsets will be easier to handle than overconvergent subanalytic subsets, and we'll be able to prove the theorems mentioned above for overconvergent constructible subsets and then deduce them for overconvergent subanalytic subsets thanks to theorem 1.31.

Overconvergent constructible subsets. Before explaining what overconvergent constructible subsets are, we would like to make an analogy with Chevalley's theorem in algebraic geometry. For simplicity, we want to interpret Chevalley's theorem in the following way. Let $\mathcal{X}$ be a (noetherian) affine scheme:

- In one hand, one can consider constructible subsets of $\mathcal{X}$ which are finite union of locally closed subsets. Since $\mathcal{X}$ is affine, the constructible subsets of $\mathcal{X}$ are the finite boolean combination of sets of the form $\{x \in \mathcal{X} \mid f(x) = 0\}$ where $f \in \mathcal{O}(\mathcal{X})$.
- On the other hand one can consider the subsets of $\mathcal{X}$ of the form $S = f(\mathcal{Y})$ where $f : \mathcal{Y} \to \mathcal{X}$ is a finite type morphism. Alternatively, it can be described as the subsets of the form $\pi(T)$ where T is a constructible subset of $\mathcal{X} \times \mathbb{A}^n$ and $\pi : \mathcal{X} \times \mathbb{A}^n \to \mathcal{X}$ is the natural projection.

Chevalley's theorem asserts precisely that these two families of subsets of $\mathcal{X}$ agree.

If we come back to our situation where X is an affinoid space, the definition we have given of an overconvergent subanalytic subset of X looks like the second class above when $\mathcal{X}$ is a scheme, in the sense that we have used projections. So that we would like to give a alternative definition of overconvergent subanalytic subset without taking any projection, but using only functions of the affinoid algebra $\mathcal{A}$ associated to X, i.e. we would like to have a definition which looks like the definition of constructible subsets of $\mathcal{X}$. One natural candidate is then the class of globally

semianalytic subsets of X. But this class is too small:
the class of overconvergent subanalytic subsets properly contains the class of globally semianalytic subsets.
One could then replace globally semianalytic subset by locally semianalytic subset, but this still doesn't work:
the class of overconvergent subanalytic subsets properly contains the class of globally semianalytic subsets (however, see theorem 3.13 for special cases where this is true).
So things are not as simple as in Chevalley's theorem, and one needs to add new functions. One typical example is the following:

Example 0.4. Let X be an affinoid space of the form $X = \mathcal{M}(\mathcal{A})$. Let $f, g \in \mathcal{A}$, $\mathbb{B} = \mathcal{M}(k\{t\})$, $T = \{z \in X \times \mathbb{B} \mid f(z) = (tg)(z)\}$ and $S = \pi(T)$ where $\pi : X \times \mathbb{B} \to X$ is the natural projection. Then $S = \{x \in X \mid |f(x)|| \leq |g(x)|\}$ is still a globally semianalytic subset of X. But if in addition we consider $h = h(t) \in \mathcal{A}\{t\}$ a series whose radius of convergence in t is > 1, and define $T' = \{z \in T \mid |h(z)| \leq 1\}$, it is not clear at all that $\pi(T')$ will be globally semianalytic. But it is overconvergent subanalytic. Although it is not semianalytic, one would like to say that

$$T' = \{z \in T \mid |h\left(\frac{f}{g}\right)(z)| \leq 1\}.$$

We now explain on this example, what is the definition of an overconvergent subanalytic subset: we set $Y = \mathcal{M}(\mathcal{B})$ whose affinoid algebra is $\mathcal{B} = \mathcal{A}\{r^{-1}t\}/(f - tg)$ where $r > 1$. There is a natural map $Y \xrightarrow{\varphi} X$ which is actually one of the two natural charts of the blowing up of X along the ideal (f, g). Now, $\{y \in Y \mid g(y) \neq 0\}$ corresponds to the analytic domain of X: $\{x \in X \mid |f(x)| \leq |g(x)| \neq 0\}$ and the function $t \in \mathcal{B}$ corresponds to $\frac{f}{g}$ on it. Hence in the example above, T' was the projection along φ of a globally semianalytic subset T of Y such that $T \subseteq \{y \in Y \mid |\frac{f}{g}(y)| \leq 1\}$. In general, we iterate this procedure: we define a constructible datum as a map $Y \overset{\varphi}{\dashrightarrow} X$ which is a composition of blowing up as above, and consider projections $\varphi(T) \subseteq X$ of globally semianalytic subsets T of Y which lives outside the exceptional locus of the blowing up φ. An overconvergent constructible subset of X is then a finite union of such sets $\varphi(T)$.
Let us now explain how is proven theorem 1.31 which asserts that overconvergent subanalytic subsets are precisely the overconvergent constructible subsets. It is quite an easy consequence of the definition that overconvergent constructible subsets are overconvergent subanalytic. To prove the reverse inclusion, we'll be in the following situation: T will be a globally semianalytic subset of $X \times \mathbb{B}^n$ defined with overconvergent functions. Then we'll algebraize these function with Weierstrass theory (proposition 1.30) and our constructible data. Finally we'll use quantifier elimination in the theory of algebraically closed valued fields.

Organisation of the paper.
In section 1, we define *constructible data* of X, in order to define overconvergent constructible subsets. Note that we don't assume that k is algebraically closed contrary to [19]. In section 1.2 we introduce overconvergent subanalytic subsets. In section 1.3 we've tried to carefully treat Weierstrass division, trying to be as general as possible (namely our results hold for an arbitrary Banach algebra, and an arbitrary radius of convergence). In section 1.4 we prove that overconvergent constructible and overconvergent subanalytic subsets are the same. The proof of this result which is in [19], is simplified by the use of Berkovich spaces: the quite technical section 2 of [19] *A combinatorial lemma* is replaced by a simple compactness argument (see theorem 1.31). In 1.5 we try to handle the following problem: how to pass from a definition that works only for affinoid spaces to a more local definition, with the hope that in the affinoid case the local and the global definitions would coincide. As we said earlier, trying to do this with the G-topology won't work. If in contrary we do this with the Berkovich topology, the definitions will be compatible. This can be summed up by:

Theorem. *Let S be a subset of the affinoid space X. The following properties are equivalent:*

1. *S is overconvergent constructible.*
2. *S is overconvergent subanalytic.*
3. *There exists $r > 1$, an integer n, T a locally semianalytic subset of $X \times \mathbb{B}^n_r$ such that $S = \pi(T \cap (X \times \mathbb{B}^n))$ where $\pi : X \times \mathbb{B}^n \to X$ is the natural projection.*

(4) S is, locally for the Berkovich topology, an overconvergent constructible subset.

(1) ⇔ (2) is theorem 1.31, (1) ⇔ (3) is corollary 1.39, and (1) ⇔ (4) is proposition 1.35.

In section 2, we give some counter-examples to erroneous statements of [19]. Precisely, in [19] five classes of subsets were defined: globally strongly subanalytic, globally strongly D-semianalytic, strongly subanalytic, locally strongly subanalytic and strongly D-semianalytic subsets. The three last classes were defined from first the two ones by adding "G-local" at some point. In [19] it was claimed that these five classes agree. In section 2 we explain that this is not the case, namely from these five classes, the two first one indeed agree, but not the three last ones, which are larger (see figure 1). What the above theorem says is that if one replaces "G-local" by "locally for the Berkovich topology", the results of [19], for instance the theorem on p. 270, becomes true.
In section 2 we then give various counterexamples. Let us explain one of them:

Example 0.5. Let $X = \mathbb{B}^2$ be the the closed bidisc, $0 < r < 1$ with $r \in \sqrt{|k^\times|}$, $f \in k\{r^{-1}x\}$ an analytic function whose radius of convergence is exactly r and such that $\|f\| < 1$. We then define

$$S = \{(x,y) \in \mathbb{B}^2 \mid |x| < r \text{ and } y = f(x)\}$$

Then (see proposition 2.4) S is rigid semianalytic, but not overconvergent subanalytic. The Berkovich approach is here helpful since to prove this, we use a point of the Berkovich bidisc (and properties of its local ring) that is not a rigid point.

Finally, in section 3 we correct the proof of [20] (which rested on the erroneous results of [19], and [21]) and restrict the hypothesis of it. Namely, we prove that when k is algebraically closed, and X is a quasi-smooth (equivalently rig-smooth) k-analytic space of dimension 2, then overconvergent subanalytic subsets are in fact locally semianalytic. Note only we give a correct proof of this theorem, but moreover, this result is more general than the result of [20], where X was the bidisc and where is was assumed that the characteristic of k was 0.

Contribution of this text. We want to stress the fact that section 1 is highly inspired by the work of H. Schoutens. In particular, the definition we give of a constructible datum, and the resulting definition of an overconvergent constructible subset, is just a *geometric* formulation of what is done in [19] concerning D-strongly semianalytic subsets. In particular, the proof of theorem 1.31 is very close to the proof of [19, Th 5.2]. We have however decided to include a proof of theorem 1.31 for three reasons. First, the compactness argument we use in theorem 1.31 seemed to us enlightening, and a way to see that Berkovich spaces are relevant in this context[1]. Secondly, we have the feeling that replacing the strongly D-semianalytic subsets of [19] by our constructible data, and overconvergent constructible subsets is a way to stay at the level of geometry and to have a better understanding of the situation. Finally, the mistakes in [19], that we explain in section 2, have the following consequences: almost all the statements of [19] become false. For instance, [19, Theorem 5.2] asserts that strongly subanalytic subsets are equivalent to globally strongly D-semianalytic subsets, which is false as we prove in section 2. The correct assertion would have been that globally strongly subanalytic subsets are equivalent to globally strongly D-semianalytic subsets. In this context it seemed to us relevant to write section 1.
The same remarks hold for section 3. An statement analogous to theorem 3.13 was claimed in [20]. However, in this article, it was assumed, and used in the proofs, that the five classes of subsets introduced in [19] were the same, so that the proofs, and maybe the statements, are erroneous.
Finally, let us mention that another proof of theorem 1.31 has been given in [5, 4.4 (10)].

1. OVERCONVERGENT CONSTRUCTIBLE SUBSETS

1.1. **Constructible data.** With a few exceptions that will be specified, X will always be an affinoid space of the form $\mathcal{M}(\mathcal{A})$.

Definition 1.1. Let X be an affinoid space whose affinoid algebra is $\mathcal{A}$. A subset S of X is called *globally semianalytic* if it is a finite boolean combination of sets of the form $\{x \in X \mid |f(x)| \leq |g(x)|\}$ where f and $g \in \mathcal{A}$ (by finite boolean combination, we mean finitely many use of the set-theoretical operators $\cap$, $\cup$ and ^{c}).

---

[1] However, it has to be noted that we could have written this proof in the context of adic spaces, and use a similar argument of quasi-compactness.

A subset of the form $\{x \in X \mid |f_i(x)| \Diamond_i |g_i(x)|\ i = 1 \dots n\}$ with f_i and $g_i \in \mathcal{A}$, and $\Diamond_i \in \{\leq, <\}$ will be called *basic semianalytic.*

With a repeated use of the rule $(A \cup B) \cap C = (A \cap C) \cup (B \cap C)$ one can show that any globally semianalytic subset of X is a finite union of basic semianalytic subsets.

Definition 1.2. Let (X, S) be a *k-germ* of k-analytic space (this just means that S is a subset of X [2, 3.4]). An elementary constructible datum of (X, S), is the following datum. Let $f, g \in \mathcal{A}$. Let also r and s be two real numbers such that $r > s > 0$ and $s \in \sqrt{|k^*|}$. Let

$$Y = \mathcal{M}(\mathcal{A}\{r^{-1}t\}/(f - tg)) \xrightarrow{\varphi} \mathcal{M}(\mathcal{A}) = X$$

and let $R \subseteq Y$ be a globally semianalytic subset of Y. Define

$$T := \varphi^{-1}(S) \cap \{y \in R \mid g(y) \neq 0 \text{ and } |f(y)| \leq s|g(y)|\}$$

Then $(Y, T) \xrightarrow{\varphi} (X, S)$ is an elementary constructible datum.
If $\psi : (Y', T') \simeq (Y, T)$ is an isomorphism of *k-germs*, and $(Y, T) \xrightarrow{\varphi} (X, S)$ is an elementary constructible datum, if we set $\varphi' = \varphi \circ \psi$, then we'll also say that $(Y', T') \xrightarrow{\varphi'} (X, S)$ is an elementary constructible datum.

If S is a globally semianalytic subset of X, one checks that if $(Y, T) \xrightarrow{\varphi} (X, S)$ is an elementary constructible datum, T is necessarily a globally semianalytic subset of Y. So in that case, instead of having defined first a globally semianalytic subset R of Y, and afterwards T, we could alternatively have defined an elementary constructible datum $(Y, T) \xrightarrow{\varphi} (X, S)$ with f, g, r, s, Y as above, and with T a globally semianalytic subset of Y such that

$$T \subseteq \varphi^{-1}(S) \cap \{y \in Y \mid g(y) \neq 0 \text{ and } |f(y)| \leq s|g(y)|\}$$

Note that we use the notion of *k-germs*, (X, S), which must not be confused with the notion of *k-$\mathscr{G}$erms* ([2, 3.4]).

Remark 1.3. If $(Y, T) \xrightarrow{\varphi} (X, S)$ is an elementary constructible datum, then $\varphi(T) \subset S$, and φ realizes a homeomorphism between T and its image $\varphi(T)$. Moreover $\{y \in Y \mid g(y) \neq 0 \textit{ and } |f(y)| \leq s|g(y)|\}$ is an analytic domain of Y, and can be identified through φ with an analytic domain of X.

Definition 1.4. Let (X, S) be a k-germ. A *constructible datum* is a sequence

$$(Y, T) = (X_n, S_n) \xrightarrow{\varphi_n} (X_{n-1}, S_{n-1}) \to \cdots \to (X_1, S_1) \xrightarrow{\varphi_1} (X_0, S_0) = (X, S)$$

where for $i = 1 \dots n$, $(X_i, S_i) \xrightarrow{\varphi_i} (X_{i-1}, S_{i-1})$ is an elementary constructible datum. Let $\varphi = \varphi_1 \circ \dots \circ \varphi_n$. Then we'll note this constructible datum $(Y, T) \overset{\varphi}{\dashrightarrow} (X, S)$. We'll say that the complexity of φ is n.

In the particular case $S = X$, i.e. $(X, S) = (X, X)$, we'll note it $(Y, T) \overset{\varphi}{\dashrightarrow} X$. This is actually the case that will mainly interest us, but partly for technical reasons we have chosen to use *k-germs*. The previous remark implies that if $(Y, T) \overset{\varphi}{\dashrightarrow} (X, S)$ is a constructible datum, $\varphi_{|T} : T \xrightarrow{\varphi_{|T}} S$ induces a homeomorphism between T and $\varphi(T)$. It is also clear that if $(Z, U) \overset{\psi}{\dashrightarrow} (Y, T)$ is a constructible datum and $(Y, T) \overset{\varphi}{\dashrightarrow} (X, S)$ is another one, then $(Z, U) \overset{\varphi \circ \psi}{\dashrightarrow} (X, S)$ is also a constructible datum.

Definition 1.5. Let $(X_i, S_i) \overset{\varphi_i}{\dashrightarrow} (X, S)$, $i = 1 \dots m$ be m constructible data of the k-germ (X, S). We'll say it forms a *constructible cover* of (X, S) if $\bigcup\limits_{i=1}^{m} \varphi_i(S_i) = S$.

Definition 1.6. Let (X, S) be a k-germ. A subset C of S is said to be an *overconvergent constructible subset* of (X, S) if there exist m constructible data $(X_i, S_i) \overset{\varphi_i}{\dashrightarrow} (X, S)$ for $i = 1 \dots m$ such that $\bigcup\limits_{i=1}^{m} \varphi_i(S_i) = C$.
If $S = X$, we'll simply say that C is an overconvergent constructible subset of X (instead of (X, X)).

Remark 1.7. Using notations of def. 1.2, when $(Y,T) \xrightarrow{\varphi} (X,S)$ is an elementary constructible datum, with $Y = \mathcal{M}(\mathcal{B})$, then T (and hence $\varphi(T)$) are defined with the function t which mimics the function $\frac{f}{g}$, when it has a sense and its norm is $\leq s$. In addition the condition $r > s$ is here to make sure that the new functions of $\mathcal{B}$, seen as functions on the domain defined by $\{x \in X \ \big| |f(x)| \leq s|g(x)|\}$ are overconvergent in t.

Lemma 1.8. *If $(Y,T) \xrightarrow{\varphi} (X,S)$ is an elementary constructible datum and $(Z,U) \xrightarrow{\varphi} (X,S)$ is a morphism of k-germs, let us consider the cartesian product of k-germs:*

$$\begin{tikzcd}
(Y,T) \arrow[r, "\varphi"] & (X,S) \\
(Y,T)\times_{(X,S)}(Z,U) \arrow[u, "\psi'"] \arrow[r, "\varphi'"] & (Z,U) \arrow[u, "\psi"]
\end{tikzcd}$$

Then, $(Y,T)\times_{(X,S)}(Z,U) \xrightarrow{\varphi'} (Z,U)$ is an elementary constructible datum. Moreover if we set

$$(Y,T)\times_{(X,S)}(Z,U) = (Y',T')$$

then, $(\varphi\circ\psi')(T') = \varphi(T)\cap\psi(U)$.

Corollary 1.9. *Let $(Y,T) \overset{\varphi}{\dashrightarrow} (X,S)$ be a constructible datum*

$$(Y,T) = (X_n,S_n) \xrightarrow{\varphi_n} \cdots \xrightarrow{\varphi_1} (X_0,S_0) = (X,S)$$

And let $(X',S') \xrightarrow{\psi} (X,S)$ be a morphism of k-germs. Let us consider the cartesian product :

$$\begin{tikzcd}
(Y,T) \arrow[r, dashed, "\varphi"] & (X,S) \\
(Y',T') \arrow[u, "\psi'"] \arrow[r, dashed, "\varphi'"] & (X',S') \arrow[u, "\psi"]
\end{tikzcd}$$

Then $(Y',T') \overset{\varphi'}{\dashrightarrow} (X',S')$ is a constructible datum and $(\psi\circ\varphi')(T') = \varphi(T)\cap\psi(S')$.

Corollary 1.10. *Let $(X_1,T_1) \overset{\varphi}{\dashrightarrow} (X,S)$ and $(X_2,T_2) \overset{\psi}{\dashrightarrow} (X,S)$ be two constructible data (with the same target). Let us consider the fibered product*

$$\begin{tikzcd}
(X_1,T_1) \arrow[r, dashed, "\varphi"] & (X,S) \\
(Z,U) \arrow[u, dashed, "\psi'"] \arrow[r, dashed, "\varphi'"] & (X_2,T_2) \arrow[u, dashed, "\psi"]
\end{tikzcd}$$

Then $(Z,U) \overset{\psi'}{\dashrightarrow} (X_1,T_1)$ and $(Z,U) \overset{\varphi'}{\dashrightarrow} (Y_2,T_2)$ are constructible data. Moreover $(\varphi\circ\psi')(U) = (\psi\circ\varphi')(U) = \varphi(T_1)\cap\psi(T_2)$.

By definition, an overconvergent constructible subset C of (X,S) is a subset of S, and when we'll say *the complementary subset of C*, this will mean $S\setminus C$.

Proposition 1.11.

1. *If $T\subseteq X$ is a globally semianalytic subset of X then $T\cap S$ is an overconvergent constructible subset of (X,S).*
2. *Let $C\subseteq T$ be an overconvergent constructible subset of (Y,T) and $(Y,T) \overset{\varphi}{\dashrightarrow} (X,S)$ a constructible datum. Then $\varphi(C)$ is on overconvergent constructible subset of (X,S).*
3. *The class of overconvergent constructible subset of (X,S) is stable under finite boolean combination.*

Proof.

(1) Consider the elementary constructible datum $(X, T\cap S) \xrightarrow{id} (X,S)$.

(2) By definition, there exist some constructible data $(Y_i,T_i) \overset{\varphi_i}{\dashrightarrow} (Y,T)$, for $i = 1 \dots m$, such that $C = \bigcup\limits_{i=1}^{m} \varphi_i(T_i)$. Now if we define $\psi_i := \varphi \circ \varphi_i$, then $(Y_i,T_i) \overset{\psi_i}{\dashrightarrow} (X,S)$ are m constructible data, and $\varphi(C) = \varphi(\bigcup\limits_{i=1}^{m} \varphi_i(Y_i)) = \bigcup\limits_{i=1}^{m} \psi_i(T_i)$, so it is an overconvergent constructible subset of (X,S).

(3) Stability under finite union is a direct consequence of the definition, as for intersection, it is a consequence of corollary 1.10. And if $C \subseteq S$ is an overconvergent constructible subset of (X,S), let's show that $S \setminus C$ is also overconvergent constructible. By definition, $C = \bigcup\limits_{i=1}^{m} \varphi_i(S_i)$ where $(X_i,S_i) \overset{\varphi_i}{\dashrightarrow} (X,S)$ are some constructible data. We do it by induction on c, the maximum of the complexity of the φ_i's.
If $c = 0$, there exists T a globally semianalytic subset of X such that $C = T \cap S$. Then $S \setminus C = S \cap (T^c)$ is then overconvergent constructible.
If $c > 0$ and we assume the result for $c' < c$, then

$$S \setminus C = S \setminus (\bigcup_{i=1}^{m} \varphi_i(S_i)) = \bigcap_{i=1}^{m} (S \setminus \varphi_i(S_i))$$

so we can assume that $m = 1$, that is to say, we can assume that $C = \varphi(T)$ where $(Y,T) \overset{\varphi}{\dashrightarrow} (X,S)$ is a constructible datum of complexity c. Then $\varphi = \psi \circ \varphi' : (Y,T) \overset{\varphi'}{\dashrightarrow} (Y',T') \xrightarrow{\psi} (X,S)$ where the complexity of φ' is $c-1$ and ψ is an elementary constructible datum. Now $S \setminus \varphi(T) = \psi(T' \setminus \varphi'(T)) \cup (S \setminus \psi(T'))$ (because $\varphi'_{|T}$ and $\psi_{|T'}$ are injective maps). By induction hypothesis, $T' \setminus \varphi'(T)$ is an overconvergent constructible subset of (Y',T'), so according to (1), so is $\psi(T' \setminus \varphi'(T))$.
Finally, if the elementary constructible datum ψ is associated with f, g, r and s, by definition, $T' = \psi^{-1}(S) \cap \{y \in R \mid |f(y)| \leq s|g(y)| \neq 0\}$ for some globally semianalytic subset R of Y'. And if we define $\tilde{T} = \psi^{-1}(S) \cap \{y \in R^c \mid |f(y)| \leq s|g(y)| \neq 0\}$, then

$$S \setminus \psi(T') = \psi(\tilde{T}) \cup \{y \in S \mid |f(y)| > s|g(y)|\} \cup \{y \in S \mid g(y) = 0\}$$

is also overconvergent constructible in (X,S).

$\square$

Let $x \in X$, and U be an affinoid neighbourhood of x. Shrinking U if necessary, we can assume [1, 2.5.15] that U is a rational domain of the form $X(\underline{r}^{-1}\frac{f}{g}) = \{p \in X \mid |f_i(x)| \leq r_i|g(x)|\}$ such that $X\left(\left(\frac{\underline{r}}{2}\right)^{-1}\frac{f}{g}\right)$ still contains x. For each i, we pick a real number s_i such that $\frac{r_i}{2} < s_i < r_i$ and $s_i \in \sqrt{|k^\times|}$. For each i, we consider the elementary constructible datum $(X_i,S_i) \xrightarrow{\varphi_i} X$ defined by $X_i = \mathcal{A}\{r_i^{-1}t_i\}/(f_i - t_i g)$, and $S_i = \{p \in X_i \mid |f_i(p)| \leq s_i|g(p)| \text{ and } g(p) \neq 0\}$. One checks that $\varphi_i(S_i)$ is a neighbourhood of x. Now if we take the fibered product of all these elementary constructible data, we obtain (using corollary 1.10) the following constructible datum:

$$\left(X\left(\underline{r}^{-1}\frac{f}{g}\right), X\left(\underline{s}^{-1}\frac{f}{g}\right)\right) \overset{\varphi}{\dashrightarrow} X$$

Here φ just corresponds to the embedding of the affinoid domain $X(\underline{r}^{-1}\frac{f}{g})$. Moreover $\varphi\left(X(\underline{s}^{-1}\frac{f}{g})\right)$, that we might identify with $X\left(\underline{s}^{-1}\frac{f}{g}\right)$, is a neighbourhood of x. We can sum up this in the following lemma:

Lemma 1.12. *Let $x \in X$ and U be an affinoid neighbourhood of x. Then there exists a constructible datum $(Y,T) \overset{\varphi}{\dashrightarrow} X$ such that T is an affinoid domain of Y, φ is the embedding of an affinoid domain $Y \to X$ such that Y is in fact an affinoid subdomain of U, and $\varphi(T)$ is an affinoid neighbourhood of x.*

1.2. **Overconvergent subanalytic subsets.** We'll denote by $\mathbb{B}$ (resp. $\mathbb{B}_r$ for $r > 0$ a real number) the closed disc of radius 1 (resp. r), and if n is an integer, $\mathbb{B}^n$ and $\mathbb{B}_r^n$ will denote the correspond closed polydiscs.
More generally, for $\underline{r} = (r_1, \dots, r_n) \in {\mathbb{R}_+^*}^n$ a polyradius, we'll denote by

$$\mathbb{B}_{\underline{r}} = \mathcal{M}(k\{\underline{r}^{-1}t\}) = \mathcal{M}(k\{r_1^{-1}t_1, \dots, r_n^{-1}t_n\})$$

the polydisc of radius $\underline{r}$, and $\overset{\circ}{\mathbb{B}}(\underline{r})$ the corresponding open polydisc. When the number n will be clear from the context, we'll write $\underline{1}$ for $(1, \dots, 1) \in \mathbb{R}^n$, and $\underline{0}$ or 0 for $(0, \dots, 0) \in \mathbb{R}^n$. Finally, $\underline{\rho} > \underline{r}$ will mean that $\rho_i > r_i$ for $i = 1 \dots n$.

Definition 1.13. A subset $S \subset X$ is said to be an overconvergent subanalytic subset of X if there exist $n \in \mathbb{N}$, $r > 1$, and $T \subseteq X \times \mathbb{B}_r^n$ a globally semianalytic subset such that $S = \pi(T \cap (X \times \mathbb{B}^n))$ where $\pi : X \times \mathbb{B}_r^n \to X$ is the natural projection.

Lemma 1.14. *Let $f : Y \to X$ be a morphism of affinoid spaces and S an overconvergent subanalytic subset of X. Then $f^{-1}(S)$ is an overconvergent subanalytic subset of X.*
In particular, if V is an affinoid domain of X and S an overconvergent subanalytic subset of X, $S \cap V$ is an overconvergent subanalytic subset of V.

Proof. Let $r > 1$ and $T \subseteq X \times \mathbb{B}_r^n$ be a globally semianalytic subset such that $S = \pi(T \cap (X \times \mathbb{B}^n))$. Let us consider the following cartesian diagram:

(1)
$$\begin{tikzcd}
Y \times \mathbb{B}_r^n \arrow[r, "f'"] \arrow[d, "\pi'"] & X \times \mathbb{B}_r^n \arrow[d, "\pi"] \\
Y \arrow[r, "f"] & X
\end{tikzcd}$$

Then $f^{-1}(S) = f^{-1}(\pi(T \cap (X \times \mathbb{B}^n))) = \pi'(f'^{-1}(T \cap (X \times \mathbb{B}^n)))$. The last inequality holds because (1) is a cartesian diagram. Now $\pi'(f'^{-1}(T \cap (X \times \mathbb{B}^n))) = \pi'(f'^{-1}(T) \cap (Y \times \mathbb{B}^n)) = \pi'^{-1}(T' \cap (Y \times \mathbb{B}^n))$ where $T' = f'^{-1}(T)$ is a globally semianalytic subset of $Y \times \mathbb{B}_r^n$. Hence $f^{-1}(S) = \pi'(T' \cap (Y \times \mathbb{B}^n))$ is an overconvergent subanalytic subset of Y. ☐

We could alternatively have defined an overconvergent subanalytic subset of X as some $\pi(T)$ where T is a globally semianalytic subset of $X \times \mathbb{B}_r^n$ such that $T \subseteq X \times \mathbb{B}^n$. This indeed gives an equivalent definition because $X \times \mathbb{B}^n$ is a globally semianalytic subset of $X \times \mathbb{B}_r^n$.

Remark 1.15. For $\underline{s} \in \sqrt{|k^\times|}^n$, $k\{\underline{s}^{-1}X\}$ is a strictly k-affinoid algebra ([1, 2.1.1] and [3, 6.1.5.4]). Hence if $\underline{r} > \underline{s}$, and $T \subseteq X \times \mathbb{B}_{\underline{r}}$ is a globally semianalytic subset, then $\pi\left(T \cap (X \times \mathbb{B}_{\underline{s}})\right)$ is an overconvergent subanalytic subset of X.
Indeed let $\underline{s} \in \sqrt{|k^\times|}^n$ and $\underline{r} \in \mathbb{R}^n$ such that $\underline{s} < \underline{r}$, and $T \subseteq X \times \mathbb{B}_{\underline{r}}$ a globally semianalytic subset of $X \times \mathbb{B}_{\underline{r}}$. Let's show that $\pi(T \cap (X \times \mathbb{B}_{\underline{s}})$ is overconvergent subanalytic in the sense of definition 1.13. First to simplify notations, we can assume that $n = 1$, i.e. that $s \in \sqrt{|k^\times|}$ and $r \in \mathbb{R}$ (the proof for an arbitrary n is the same). Then, up to a multiplication by some $\mu \in k^\times$ small enough, we can assume that $s \leq 1$. Since $s \in \sqrt{|k^\times|}$, there exist $\lambda \in k^\times$ and $m \in \mathbb{N}$ such that $s^m = |\lambda|$. Then in

$$\mathbb{B}_{(r, (\frac{r}{s})^m)} = \mathcal{M}(k\{r^{-1}y, \left(\left(\frac{r}{s}\right)^m\right)^{-1} t\})$$

let's consider the Zariski-closed subset defined by $y^m = \lambda t$, i.e. $V(y^m - \lambda t)$. Then, the map:

$$\begin{array}{ccc} \mathbb{B}_r & \to & \mathbb{B}_{(r, (\frac{r}{s})^m)} \\ x & \mapsto & (x, \frac{x^m}{\lambda}) \end{array}$$

identifies $\mathbb{B}_r$ with the Zariski closed subset $V(y^m - \lambda t)$ and moreover, since $s \leq 1$

$$\begin{array}{ccc} \mathbb{B}_s & \to & \mathbb{B}^2 \\ x & \mapsto & (x, \frac{x^m}{\lambda}) \end{array}$$

identifies $\mathbb{B}_s$ with the Zariski-closed subset of $\mathbb{B}^2$, $V(y^m - \lambda t)$. Taking the fibre product with X we then obtain:

$$\begin{tikzcd}
X \times \mathbb{B}_r \arrow[r, "\simeq"] & V(y^m - \lambda t) \arrow[r, hook, "\alpha"] & X \times \mathbb{B}_{(r,(\frac{r}{s})^m)} \\
X \times \mathbb{B}_s \arrow[u, hook] \arrow[r, "\simeq"] \arrow[dr, "\pi"'] & V(y^m - \lambda t) \arrow[u, hook] \arrow[r, hook, "\beta"] & X \times \mathbb{B}^2 \arrow[u, hook] \arrow[dl, "\pi"] \\
 & X &
\end{tikzcd}$$

Hence if $T \subseteq X \times \mathbb{B}_r$ is globally semianalytic, $T' := \alpha(T)$ is also globally semianalytic in $X \times \mathbb{B}_{(r,(\frac{r}{s})^m)}$ and $\alpha(T) \cap (X \times \mathbb{B}^2) = \beta(T \cap (X \times \mathbb{B}_s))$. So $\pi(T \cap (X \times \mathbb{B}_s)) = \pi(T' \cap (X \times \mathbb{B}^2))$ is well overconvergent subanalytic in the sense of definition 1.13.

We now take a few lines to explain an alternative definition which emphasizes the relevance of k-$\mathscr{G}$*erms* in this context.

Definition 1.16. Let (X, S) be a k-$\mathscr{G}$*erm* [2, 3.4]. A subset $T \subseteq S$ is a globally semianalytic subset of (X, S) if there exists a representative (W, S) of (X, S), with W an affinoid space, and $R \subseteq W$ a globally semianalytic subset of W such that $T = S \cap R$.

Lemma 1.17. *Let* $\mathcal{W}$ *be a neighbourhood of* $X \times \mathbb{B}^n$ *in* $X \times \mathbb{A}_k^{n,an}$. *Then there exists* $r > 1$ *such that* $\mathcal{W} \supseteq X \times \mathbb{B}_r^n$.

Proof. If necessary, we can assume that $\mathcal{W} \subseteq X \times \mathbb{B}_s^n$ with $s > 1$ and also assume that $\mathcal{W}$ is open. In that case, $Z := (X \times \mathbb{B}_s^n) \backslash \mathcal{W}$ is a compact subset of $X \times \mathbb{B}_s^n$, and by assumption $Z \cap (X \times \mathbb{B}^n) = \emptyset$, so

$$Z \subseteq \bigcup_{i=1}^{n} \bigcup_{r>1} \{p \in X \times \mathbb{B}_s^n \bigm| |T_i(p)| > r\}$$

Hence by compactness there exists $r > 1$ such that $Z \subseteq \bigcup\limits_{i=1}^{n} \{p \in X \times \mathbb{B}_s^n \bigm| |T_i(p)| > r_i\}$, which says that $\mathcal{W} \supseteq X \times \mathbb{B}_r^n$. ☐

Hence S is an overconvergent subanalytic subset of X if and only if there exist an integer n, and T a semianalytic subset of the k-$\mathscr{G}$*erm* $(X \times \mathbb{A}_k^{n,an}, X \times \mathbb{B}^n)$ such that $S = \pi(T)$.

1.3. **Weierstrass preparation.** In this section, A will be a (ultrametric) complete normed ring i.e. satisfies the inequality $\|ab\| \leq \|a\|.\|b\|$ [3, 1.2.1.1].
If $r > 0$, the norm $\|.\|$ that we'll consider on $A\{r^{-1}X\}$ will always be the following one: if $g = \sum\limits_{n \in \mathbb{N}} a_n X^n \in A\{r^{-1}X\}$ then $\|g\| = \max\limits_{n \geq 0} \|a_n\| r^n$.

If $m \in \mathbb{N}$, we'll note $A_m[X]$ the subset of $A[X]$ made of the polynomials of degree less or equal to m.

Definition 1.18. An element $u \in A$ is a multiplicative unit if u is invertible and for all $a \in A$, $\|u.a\| = \|u\|.\|a\|$.

Note that if u and v are multiplicative units, so is uv.

Lemma 1.19. *An element* $u \in A^*$ *is a multiplicative unit if and only if* $u \in A^*$ *and* $\|u^{-1}\| = \|u\|^{-1}$.

Proof. If u is a multiplicative unit, $1 = \|uu^{-1}\| = \|u\|\|u^{-1}\|$, so $\|u^{-1}\| = \|u\|^{-1}$.
Conversely if u is invertible and $\|u^{-1}\| = \|u\|^{-1}$, then if $a \in A$, $\|a\| = \|u^{-1}(ua)\| \leq \|u^{-1}\|\|ua\| = \|u\|^{-1}\|ua\|$. So $\|ua\| \geq \|u\|\|a\|$. Since in any case the reverse inequality $\|ua\| \leq \|u\|\|a\|$ holds, we conclude that $\|ua\| = \|u\|\|a\|$. ☐

Remark 1.20. From that, we get that if $u \in A$ and $\|u\| < 1$, then $(1 + u)$ is a multiplicative unit because

$$\|1 + u\| = 1 = \|\sum_{n \geq 0} (-u)^n\| = \|(1 + u)^{-1}\|$$

Let us note also that if u is a multiplicative unit, for all $x \in \mathcal{M}(A)$, $|u(x)| = \|u\|$. Indeed, the definition of $\mathcal{M}(\mathcal{A})$ implies that $|u(x)| \leq \|u\|$, hence $1 = |u(x)||u^{-1}(x)| \leq \|u\|\|u^{-1}\| = 1$. So the inequality was in fact the equality $|u(x)| = \|u\|$.

Remark 1.21. If $\varphi : A \to B$ is a contractive morphism of normed ring (i.e. $\|\varphi(a)\| \leq \|a\|$ for all a in A), then φ sends multiplicative units to multiplicative units.
Indeed we have the sequence of inequalities:

$$1 = \|\varphi(u)\varphi(u)^{-1}\| \leq \|\varphi(u)\|\|\varphi(u^{-1})\| \leq \|u\|\|u^{-1}\| = 1$$

So there were only equalities and $\varphi(u)$ is a multiplicative unit because $\|\varphi(u)\| = \|u\|$, and $\|\varphi(u)^{-1}\| = \|u^{-1}\| = \|u\|^{-1} = \|\varphi(u)\|^{-1}$.
This remark will apply in the following context: when $\mathcal{A}$ is a k-affinoid algebra and we look at a morphism $\varphi : \mathcal{A} \to \mathcal{B} = \mathcal{A}\{r^{-1}X\}/I$ with I any ideal, and $\mathcal{B}$ is equipped with the quotient norm inherited from $\mathcal{A}\{r^{-1}X\}$. In this situation, φ is contractive. This is the case when we consider φ the morphism of a constructible datum $(Y,T) \overset{\varphi}{\dashrightarrow} (X,S)$.
Note that if φ is not contractive, multiplicative units are not necessarily preserved. For instance consider $\mathcal{A} = k\{t\}$ and $\mathcal{B} = k\{2^{-1}x, y\}/(y - x^2)$ that we equip with the residue norm. These affinoid algebras are isomorphic through $\varphi : t \mapsto x$, and if we choose $\pi \in k$ such that $2|\pi| > 1$ and $|\pi| < 1$, then $u := 1 + \pi t$ is a multiplicative unit of $\mathcal{A}$, but not $\varphi(u)$. Note however that if the field k is stable (in our situation, where k is a non-archimedean complete field, for instance k is stable if it is algebraically closed, a finite extension of a discrete valued field, or $char(\tilde{k}) = 0$ [3, 3.6.2]), one might say that any morphism of reduced affinoid algebra is contractive. Indeed, if k is stable, and $\mathcal{A}$ is a reduced affinoid algebra, then it is a distinguished affinoid algebra [3, 6.4.3], i.e. , the supremum seminorm is a residue norm on $\mathcal{A}$. If $\mathcal{B}$ is also reduced, so that the supremum seminorm is an admissible norm on it, and $\varphi : \mathcal{A} \to \mathcal{B}$ is a morphism of affinoid algebras, than φ is contractive.

Definition 1.22. Let $r > 0$ be a real number. An element $g = \sum_{n\geq 0} g_n X^n$ of $A\{r^{-1}X\}$ is X-distinguished of order s if g_s is a multiplicative unit, $\|g_s X^s\| = \|g\|$ and for all $n > s$, $\|g_n X^n\| = \|g_n\| r^n < \|g_s X^s\| = \|g_s\| r^s$. Note that in that case, g is necessarily a non zero element.

Remark 1.23. We can extend the previous remark saying that if $\varphi : A \to B$ is a contractive morphism and $g = \sum_{n\in\mathbb{N}} g_n X^n \in A\{r^{-1}X\}$ is X-distinguished of order s, then $\varphi(g) = \sum_{n\in\mathbb{N}} \varphi(g_n) X^n \in B\{r^{-1}X\}$ and it is an X-distinguished element of $B\{r^{-1}X\}$ of order s. This applies in particular when φ is the morphism of a constructible datum $(Y,T) \overset{\varphi}{\dashrightarrow} (X,S)$.

Lemma 1.24. *Let* $g = \sum_{m\in\mathbb{N}} g_m X^m \in A\{r^{-1}X\}$ *be* X*-distinguished of order* s*.*

1. *Then for all* $q = \sum_{k\in\mathbb{N}} q_k X^k \in A\{r^{-1}X\}$, $\|g.q\| = \|g\|\|q\|$.
2. *If we note* $gq = \sum_{l\in\mathbb{N}} c_l X^l$, *and if we assume that* $q \neq 0$, *and if we note* k_0 *the greatest rank such that* $\|q_{k_0}\| r^{k_0} = \|q\|$, *then* $\|gq\| = \|c_{s+k_0}\| r^{s+k_0}$ *and* $\|c_{s+k_0}\| = \|g_s\|\|q_{k_0}\|$.

Proof. First without any hypothesis it is true that $\|gq\| \leq \|g\|\|q\|$.
Conversely, $c_{s+k_0} = \sum_{m+k=s+k_0} g_m q_k$. Let then m, k be two integers such that $m + k = s + k_0$.
If $k > k_0$, $\|q_k\| r^k < \|q_{k_0}\| r^{k_0}$. So

$$\|g_m q_k\| r^{s+k_0} = \|g_m q_k\| r^{m+k} \leq \|g_m\| r^m \|q_k\| r^k < \|g_s\| r^s \|q_{k_0}\| r^{k_0} = \|g_s q_{k_0}\| r^{s+k_0}.$$

So $\|g_m q_k\| < \|g_s q_{k_0}\|$.
If $k < k_0$, then $m > s$, and since $\|g_m\| r^m < \|g_s\| r^s$ (because g is X-distinguished of order s), we obtain with the same reasoning, that $\|g_m q_k\| r^{s+k_0} < \|g_s q_{k_0}\| r^{s+k_0}$.
Finally, we get that $\|c_{s+k_0}\| = \|g_s q_{k_0}\| = \|g_s\|\|q_{k_0}\|$ (because g_s is a multiplicative unit), so $\|gq\| \geq \|g_s\| r^s \|q_{k_0}\| r^{k_0} = \|q\|\|g\|$. □

Proposition 1.25 (Weierstrass Division)**.** *Let* $g \in A\{r^{-1}X\}$ *be* X*-distinguished of order* s*. Then if* $f = \sum_{n\in\mathbb{N}} f_n X^n \in A\{r^{-1}X\}$ *there exists an unique couple* $(q, R) \in A\{r^{-1}X\} \times A_{s-1}[X]$ *such that* $f = gq + R$*. Moreover* $\|f\| = \max(\|g\|\|q\|, \|R\|)$*.*

Proof. In a first time, let us check that if such a couple exists, it must satisfy the equality $\|f\| = \max(\|g\|\|q\|, \|R\|)$.
First because of the ultrametric inequality, $\|f\| \leq \max(\|g\|\|q\|, \|R\|)$.
To show the reverse inequality, we distinguish two cases.
If $\|gq\| \neq \|R\|$, then $\|f\| = \max(\|gq\|, \|R\|) = \max(\|g\|\|q\|, R\|\|)$ according to lemma 1.24.
Otherwise $\|gq\| = \|g\|\|q\| = \|R\|$, and we use again lemma 1.24 and its notations (so $gq = \sum\limits_{l\in\mathbb{N}} c_l X^l$).
We get $\|gq\| = \|c_{s+k_0}\|r^{s+k_0}$. Since R is a polynomial of degree d which is strictly smaller than s, and since $f = gq+R$, and $d < s+k_0$, the coefficient f_{s+k_0} of f is c_{s+k_0}, hence $\|f\| \geq \|c_{s+k_0}\|r^{s+k_0} = \|g\|\|q\|$.
This finally proves that $\|f\| = \max(\|g\|\|q\|, \|R\|)$.
From this we can conclude that the couple (q, R) is unique because if $f = gq' + R'$ is another decomposition, we have $0 = g(q - q') + (R - R')$ and since $\|g\| \neq 0$, $\|q - q'\| = \|R - R'\| = 0$, i.e. $R = R'$ and $q = q'$.
Let us now show the existence of such a decomposition. Let

$$g' := \sum_{m=0}^{s} g_m X^m$$

In particular, $\|g\| = \|g'\|$ because g is X-distinguished of degree s. Let's note

$$\kappa := \frac{\max\limits_{m>s}(\|g_m\|)r^m}{\|g_s\|r^s} = \frac{\max\limits_{m>s}(\|g_m X^m\|)}{\|g\|}$$

Since g is X-distinguished of order s, $\kappa < 1$. Actually, if $\kappa = 0$, replace κ by $\frac{1}{2}$ for instance (any value of the interval $]0, 1[$ would work in fact in that case). In any case $\|g - g'\| \leq \kappa\|g\|$ and $\kappa \in]0, 1[$.
Lastly, let $N \in \mathbb{N}$ big enough so that

$$f' := \sum_{k=0}^{N} f_k X^k$$

fulfils $\|f - f'\| \leq \kappa\|f\|$. In particular, $\|f'\| = \|f\|$.
By definition and hypothesis, $g' \in A[X]$ is of degree s and possesses an invertible dominant coefficient, which is g_s. Hence in $A[X]$, one can carry out euclidean division by g' [15, 4.1.1], which gives $f' = g'q + R$, with $R \in A_{s-1}[X]$ and $q \in A[X]$. We can then apply the norm equality we have just shown in the first part of the proof, (because g' is also X-distinguished of order s): $\|f'\| = \max(\|g'\|\|q\|, \|R\|)$.
In particular $\|q\| \leq \frac{\|f'\|}{\|g'\|} = \frac{\|f\|}{\|g\|}$ so that

$$\|g\|\|q\| \leq \|f\|.$$

Moreover $\|R\| \leq \|f'\| = \|f\|$.
Moreover we have that

$$f = f' + (f - f') = g'q + R + (f - f') = gq + R + (f - f') + (g' - g)q$$

By definition of g' and of κ, $\|g' - g\| \leq \kappa\|g\|$, so

$$\|(g' - g)q\| \leq \|g\|\|q\|\kappa \leq \kappa\|f\| \tag{2}$$

Moreover by hypothesis

$$\|f - f'\| \leq \kappa\|f\|. \tag{3}$$

Hence if we set

$$h := f - f' + (g' - g)q = f - (gq + R)$$

then with (2) and (3) we obtain that $\|h\| \leq \kappa\|f\|$.
This construction is not what we wanted, but it allows to define by induction two Cauchy sequences $(q^i) \in A\{r^{-1}X\}$ and $(R^i) \in A_{s-1}[X]$ so that $\|f - (gq^i + R^i)\| \leq \kappa^i\|f\|$ in the following way:
we start with $(q^0, R^0) = (0, 0)$.
In order to perform the induction step, let $i > 0$ be given and let us assume that (q^i, R^i) is defined. We set $h^i := f - (gq^i + R^i)$, which by induction hypothesis fulfils $\|h^i\| \leq \kappa^i\|f\|$. According to what we have just said, we can define $q \in A\{r^{-1}X\}$ and $R \in A_{s-1}[X]$ such that $h^i = gq + R + h$

with $\|q\| \leq \frac{\|h^i\|}{\|g\|} \leq \kappa^i \frac{\|f\|}{\|g\|}$, and $\|R\| \leq \|h^i\| \leq \kappa^i \|f\|$ and $\|h\| \leq \kappa \|h^i\| \leq \kappa^{i+1}\|f\|$. Then we set $q^{i+1} := q^i + q$ and $R^{i+1} := R^i + R$.
Then $\|f - (gq^{i+1} + R_{i+1})\| = \|h^i - (gq + R)\| = \|h\| \leq \kappa^{i+1}\|f\|$. By construction $\|q^{i+1} - q^i\| = \|q\| \leq \kappa^i \frac{\|f\|}{\|g\|}$ and $\|R^{i+1} - R^i\| = \|R\| \leq \kappa^i \|f\|$, so these sequences are well Cauchy sequences. This ends our induction.
Now, by completeness of $A\{r^{-1}X\}$ and $A_{s-1}[X]$ the sequences (q^i) and (R^i) have a limit, that we denote by $q_\infty \in A\{r^{-1}X\}$ and $R_\infty \in A_{s-1}[X]$, which satisfy $f = gq_\infty + R_\infty$ as we wanted. Note that actually in our construction, $q^i \in A[X]$. ☐

Corollary 1.26. *[Weierstrass Preparation] Let $g \in A\{r^{-1}X\}$ be a X-distinguished element of order s. There exists an unique couple $(w, e) \in A_s[X] \times A\{r^{-1}X\}$ such that w is a unitary polynomial of degree s, e is a multiplicative unit, and $g = ew$.*

Proof. Using Weierstrass division, we can write $X^s = gq + R$ with $\|X^s\| = \max(\|g\|\|q\|, \|R\|)$, and $R \in A[X]_{s-1}$. Let's set

$$w := X^s - R.$$

So $w \in A_s[X]$ is a unitary polynomial, and since $X^s = gq + R$,

$$w = gq = X^s - R.$$

Since g is X-distinguished of order s, according to lemma 1.24, and if we note k_0 the greatest index such that $\|q_{k_0}\| r^{k_0} = \|q\|$, and $w = \sum_{l=0}^{s} w_l X^l$, we obtain

$$\|w\| = \|gq\| = \|(gq)_{s+k_0} X^{s+k_0}\| = \|w_{s+k_0} X^{s+k_0}\|.$$

But since $w \in A_s[X]$, necessarily, $s + k_0 = s$ and $k_0 = 0$. Hence, by definition of k_0, for all $k > 0$, $\|q_0\| > \|q_k\| r^k$.
The coefficient of degree s in gq being 1, (because $gq = w$), we have the equality

$$1 = g_0 q_s + g_1 q_{s-1} + \ldots + g_s q_0$$

and since $k_0 = 0$, and g is X-distinguished of degree s and thanks to lemma 1.24, we get that $\|g_s q_0\| > \|g_{s-i} q_i\|$ for $i = 1 \ldots s$, so $\|g_s q_0\| = \|1\| = 1$, and $g_s q_0 = 1 - (g_{s-1}q_1 + \ldots g_0 q_s)$, with $\|g_s q_1 + \ldots g_0 q_s\| < 1$. Hence $g_s q_0$ is a multiplicative unit, hence so is q_0, and $\|q_0\| = \|g_s\|^{-1}$. Hence

$$q = q_0\left(1 + \frac{q_1}{q_0} X + \ldots + \frac{q_k}{q_0} X^k + \ldots\right)$$

and since $k_0 = 0$ and q_0 is a multiplicative unit, $\|\frac{q_i}{q_0}\| r^i < 1$ for all $i > 0$. Hence

$$1 + \frac{q_1}{q_0} X + \ldots + \frac{q_k}{q_0} X^k + \ldots$$

is a multiplicative unit of $A\{r^{-1}X\}$, and q too. So $g = q^{-1}(X^s - R)$, with q^{-1} a multiplicative unit and $X^s - R$ a unitary polynomial of degree s. So if we set $e := q^{-1}$, we have the expected result.
As for the uniqueness of this decomposition, if $g = ew$, e and w being as in the statement of the corollary, then $w = X^s + R$ with $R \in A_{s-1}[X]$, and $X^s = w - R = e^{-1}g + (-R)$ which is the Weierstrass division of X^s by g. Hence e and R are unique and w too because $w = X^s + R$. ☐

When $\mathcal{A}$ is an affinoid algebra, $(r_1, \ldots, r_n)$ is a polyradius, and $A := \mathcal{A}\{r_1^{-1}X_1, \ldots, r_{n-1}^{-1}X_{n-1}\}$, if we note $r = r_n$, then $\mathcal{A}\{r_1^{-1}X_1, \ldots, r_n^{-1}X_n\} = A\{r^{-1}X\}$, and we can introduce the notion of an element X-distinguished, apply Weierstrass theory to them, which corresponds to the classical one, especially if $\mathcal{A} = k$, where we find the classical Tate algebra $k\{r_1^{-1}X_1, \ldots, r_n^{-1}X_n\}$.

Lemma 1.27. *Let $\varepsilon > 0$ be given and $\underline{r} > 0$ be a polyradius. Assume that A is noetherian, and let*

$$f = \sum_{\nu \in \mathbb{N}^n} f_\nu X^\nu \in A\{\underline{r}^{-1}X\}.$$

Then there exists a finite subset $J \subseteq \mathbb{N}^n$*, and for all* $\nu \in J$*, a series* $\phi_\nu \in A\{\underline{r}^{-1}X\}$ *satisfying* $\|\phi_\nu\| < \varepsilon$*, such that*

$$f = \sum_{\nu \in J} f_\nu (X^\nu + \phi_\nu)$$

and such that in the ϕ_ν*'s, not terms* X^μ *with* $\mu \in J$ *appear.*

Proof. Let's note $\mathcal{I}$ the ideal generated by the family $\{f_\nu\}_{\nu \in \mathbb{N}^n}$. Since A is noetherian, there exists J a finite subset of $\mathbb{N}^n$ such that $\mathcal{I} = A.(f_\nu)_{\nu \in J}$. So for all $\mu \in \mathbb{N}^n \setminus J$ one can find a decomposition $f_\mu = \sum\limits_{\nu \in J} f_\nu . a^\nu_\mu$ with $a^\nu_\mu \in A$. In fact, using [3, 3.7.3], we can even assume that there exists a real constant $C > 0$ such that for all $\mu \in \mathbb{N}^n$ and $\nu \in J$, $\|a^\nu_\mu\| \leq C\|f_\mu\|$ [2].
Then, let's define for $\nu \in J$

$$\phi_\nu = \sum_{\mu \in \mathbb{N}^n \setminus J} a^\nu_\mu X^\mu$$

Since $\|a^\nu_\mu\| \leq C\|f_\mu\|$, $\phi_\nu \in \mathcal{A}\{\underline{r}^{-1}X\}$. Hence, in $A\{\underline{r}^{-1}X\}$, the following equality is satisfied:

$$f = \sum_{\nu \in J} f_\nu (X^\nu + \sum_{\mu \notin J} a^\nu_\mu X^\mu) = \sum_{\nu \in J} f_\nu (X^\nu + \phi_\nu) \tag{4}$$

We are now in the following situation :

$$f = \sum_{\nu \in J} f_\nu (X^\nu + \varphi_\nu)$$

where J is a finite subset of $\mathbb{N}^n$, for all $\nu \in J$, $\varphi_\nu \in A\{\underline{r}^{-1}X\}$ and $\varphi_\nu = \sum_\mu a^\nu_\mu X^\mu$ with $a^\nu_\mu = 0$ for $\mu \in J$ and $\|a^\nu_\mu\| \leq C\|f_\mu\|$.
Now, if $\nu_0 \notin J$ we set $J' = J \cup \{\nu_0\}$, $\varphi'_{\nu_0} := 0$, and for $\nu \in J$, $\varphi'_\nu := \sum_\mu {a'}^\nu_\mu X^\mu$ where

$$\begin{cases} {a'}^\nu_\mu = a^\nu_\mu & \text{if } \mu \neq \nu_0 \\ {a'}^\nu_\mu = 0 & \text{if } \mu = \nu_0 \end{cases}$$

One checks that the properties mentioned above still hold, namely $\|{a'}^\mu_\nu\| \leq C\|f_\mu\|$, where the constant C has not been changed,

$$f = \sum_{\nu \in J'} f_\nu (X^\nu + \varphi'_\nu)$$

Moreover, when $|\mu| \to +\infty$, $C\|f_\mu\|\underline{r}^\mu \to 0$, so $\|a^\nu_\mu\| \xrightarrow[|\mu| \to +\infty]{} 0$ uniformly in ν. Hence if we increase J adding a finite number of elements in the way described just above, we'll manage to obtain a decomposition

$$f = \sum_{\nu \in J} f_\nu (X^\nu + \varphi_n)$$

such that $\|\varphi_\nu\| < \varepsilon$ for all $\nu \in J$. ☐

1.4. **Equivalence of the two notions.** From now on, $\mathcal{A}$ will be an affinoid algebra, and $\underline{r} \in {\mathbb{R}^*_+}^n$ a ployradius such that $\underline{r} > \underline{1}$ and we'll set $\mathcal{A}\{\underline{r}^{-1}X\} = \mathcal{A}\{r_1^{-1}X_1, \ldots, r_n^{-1}X_n\}$.
If $\nu = (\nu_1, \ldots, \nu_n) \in \mathbb{N}^n$, we'll set $|\nu|_\infty = \max\limits_{i=1\ldots n} \nu_i$.
If $\underline{r} \in {\mathbb{R}^*_+}^n$ and $\nu \in \mathbb{N}^n$, we'll set $\underline{r}^\nu = \prod_{i=1}^n r_i^{\nu_i}$.
When $\mu, \nu \in \mathbb{N}^n$, we'll say that $\mu <_{lex} \nu$ when μ is smaller than ν with respect to the lexicographic order, that is to say when there exists an index m such that $\mu_m < \nu_m$ and $\mu_{m+1} = \nu_{m+1}, \ldots, \mu_n = \nu_n$.

Since $\mathcal{A}$ is noetherian, we can apply lemma 1.27 to it.

---

[2] Indeed, consider

$$\begin{array}{cccc} \psi : & A^J & \to & \mathcal{I} \\ & (a_\nu & \to & \sum a_\nu f_\nu \end{array}$$

According to [3, 3.7.3.1], $\mathcal{I}$ is a complete normed A-module, and ψ is a continuous map of normed A-modules. Hence there exists a constant C such that $\|\psi(x)\| \leq C\|x\|$ for all $x \in A^J$.

Proposition 1.28. *Let* $f = \sum\limits_{\nu\in\mathbb{N}^n} f_\nu X^\nu \in \mathcal{A}\{\underline{r}^{-1}X\}$. *There exists a constructible covering of* X, $(X_i, S_i) \overset{\varphi_i}{\dashrightarrow} X$, $i = 1..N$, *such that, if we consider the following cartesian diagrams:*

$$\begin{tikzcd}
(X_i, S_i) \arrow[r, dashed, "\varphi_i"] & X \\
X_i \times \mathbb{B}_{\underline{r}} \arrow[u, "\pi_i"] \arrow[r, dashed, "\varphi'_i"] & X \times \mathbb{B}_{\underline{r}} \arrow[u, "\pi"]
\end{tikzcd}$$

and if we note $\mathcal{A}_i$ *the affinoid algebra of* X_i, *for all* $i = 1..N$, *there exist* $a_i \in \mathcal{A}_i$ *and* $g_i = \sum\limits_{\nu\in\mathbb{N}^n} g_{i,\nu} X^\nu \in \mathcal{A}_i\{\underline{r}^{-1}X\}$ *such that*

- *For all* i, *the family* $\{g_{i,\nu}\}_{\nu\in\mathbb{N}^n}$ *generates the unit ideal in* $\mathcal{A}_i$.
- $\varphi_i'^*(f)_{|\pi_i^{-1}(S_i)} = (a_i g_i)_{|\pi_i^{-1}(S_i)}$.

Proof. According to lemma 1.27 (here we won't use the extra condition $\|\varphi_\nu\| < \varepsilon$ of this lemma), we can find a finite subset $J \subseteq \mathbb{N}^n$, for $\nu \in J$ some $\phi_\nu \in \mathcal{A}\{\underline{r}^{-1}X\}$ and a decomposition of f:

$$f = \sum_{\nu\in J} f_\nu (X^\nu + \phi_\nu)$$

Let's fix any $r > 1$, and for each $\nu \in J$, let's consider the constructible data $(X_\nu, T_\nu) \overset{\varphi_\nu}{\dashrightarrow} X$ where the affinoid algebra of X_ν is $\mathcal{A}\{r^{-1}t_\mu\}_{\mu\in J\setminus\{\nu\}}/(f_\mu - t_\mu f_\nu)$, and

$$T_\nu = \{x \in X_\nu \mid |f_\kappa(x)| \leq |f_\nu(x)| \ \ \forall\kappa \in J \setminus \{\nu\} \text{ and } f_\nu(x) \neq 0\}$$

This gives rise to the following cartesian diagrams:

$$\begin{tikzcd}
(X_\nu, T_\nu) \arrow[r, dashed, "\varphi_\nu"] & X \\
X_\nu \times \mathbb{B}_{\underline{r}} \arrow[u, "\pi'"] \arrow[r, dashed, "\varphi'_\nu"] & X \times \mathbb{B}_{\underline{r}} \arrow[u, "\pi"]
\end{tikzcd}$$

Now,

$$\varphi_\nu'^*(f) = f_\nu(X^\nu + \phi_\nu + \sum_{\mu\in J\setminus\{\nu\}} t_\mu(X^\mu + \phi_\mu))$$

For $\nu \in J$, we set

$$g_\nu = X^\nu + \phi_\nu + \sum_{\mu\in J\setminus\{\nu\}} (t_\mu(X^\mu + \phi_\mu).$$

Hence,

$$\varphi_\nu'^*(f) = f_\nu g_\nu.$$

Moreover, if we note $g_\nu = \sum\limits_{\mu\in\mathbb{N}^n} g_{\nu,\mu} X^\mu$, the coefficient of index ν, $g_{\nu,\nu}$, is 1, so the coefficients of g_ν generate the unit ideal. Finally, let's remark that

$$\bigcup_{\nu\in J} \phi_\nu(T_\nu) = \{x \in X \mid \exists\nu \text{ such that } f_\nu(x) \neq 0\} = X \setminus V(\mathcal{I})$$

If we note $T = V(\mathcal{I})$, then $(X, T) \xrightarrow{id} X$ is an elementary constructible datum and $id^*(f)_{|T} = f_{|T} = 0 = 0.1$ Now if we regroup the constructible data $(X_\nu, T_\nu) \overset{\varphi_\nu}{\dashrightarrow} X$, for $\nu \in J$, with $(X, T) \xrightarrow{\varphi} X$, we obtain the desired constructible cover. ☐

Definition 1.29. Let $\underline{r} \in (\mathbb{R}_+^*)^n$ be a polyradius and $d_1, \ldots, d_{n-1}$ some integers such that $r_n^{d_i} \leq r_i$ for $i = 1 \ldots n-1$. Then

$$\sigma : \begin{cases} X_i \mapsto & X_i + X_n^{d_i} \qquad \text{if } i \leq n-1 \\ X_n \mapsto & X_n \end{cases}$$

is an automorphism of $\mathcal{A}\{\underline{r}^{-1}X\}$. We'll call such an automorphism (as well as the automorphism it induces on k-analytic spaces) a Weierstrass automorphism.

Proposition 1.30. *Let $\underline{r} > \underline{1}$ be a polyradius. For $i = 1 \dots m$ let $f_i = \sum\limits_{\nu \in \mathbb{N}^n} f_{i,\nu} X^\nu \in \mathcal{A}\{\underline{r}^{-1}X\}$ such that for all i, $\{f_{i,\nu}\}_{\nu \in \mathbb{N}^n}$ generates the unit ideal. Let $x \in X$. Then there exists a constructible datum $(Y,T) \overset{\varphi}{\dashrightarrow} X$, with $Y = \mathcal{M}(\mathcal{B})$, such that $\varphi(T)$ is a neighbourhood of x, and that there exists a polyradius $\underline{1} < \underline{\rho} \leq \underline{r}$, a Weierstrass automorphism σ of $Y \times \mathbb{B}_{\underline{\rho}}$*

$$\begin{tikzcd}
 & (Y,T) \arrow[r, dashed, "\varphi"] & X \\
Y \times \mathbb{B}_{\underline{\rho}} \arrow[ur, "\pi'"] \arrow[r, "\sigma", "\simeq"'] & Y \times \mathbb{B}_{\underline{\rho}} \arrow[u, "\pi'"] \arrow[r, dashed, "\varphi'"] & X \times \mathbb{B}_{\underline{\rho}} \arrow[u, "\pi"']
\end{tikzcd}$$

such that if $\psi := \varphi' \circ \sigma$, for all $i = 1 \dots m$, there exist β_i in $\mathcal{B}$ and a X_n-distinguished element $g_i \in \mathcal{B}\{\underline{\rho}^{-1}X\}$ such that $\psi^*(f_i)_{|\pi'^{-1}(T)} = (\beta_i g_i)_{|\pi'^{-1}(T)}$.

Proof. For $i \in \{1 \dots m\}$ let ν^i be the greatest index in the lexicographic order in $\mathbb{N}^n$ such that

$$\max_{\nu \in \mathbb{N}^n} |f_{i,\nu}(x)| = |f_{i,\nu^i}(x)|$$

By hypothesis the family $\{f_{i,\nu}\}_{\nu \in \mathbb{N}^n}$ generates the unit ideal so for each i, $f_{i,\nu^i}(x) \neq 0$. Let $d \in \mathbb{N}$ be an integer that we'll determine later on,

$$R := \min_{i=1\dots n} r_i$$

so $R > 1$. If s is a real number (that we'll also fix subsequently) such that

$$1 < s^{d^{n-1}} \leq R \tag{5}$$

we note

$$\underline{\rho} = (s^{d^{n-1}}, s^{d^{n-2}}, \dots, s^d, s) \in \mathbb{R}^n$$

So $\underline{\rho} > \underline{1}$. It is then easy to check that

$$\sigma : \left\{ \begin{array}{ccc}
X_1 & \mapsto & X_1 + X_n^{d^{n-1}} \\
\vdots & & \vdots \\
X_i & \mapsto & X_i + X_n^{d^{n-i}} \\
\vdots & & \vdots \\
X_{n-1} & \mapsto & X_{n-1} + X_n^d \\
X_n & \mapsto & X_n
\end{array} \right.$$

is Weierstrass automorphism of $\mathcal{A}\{\underline{\rho}^{-1}X\}$.
Let's fix temporarily $i \in \{1..m\}$. If (to simplify notations) we write $f_i = f = \sum\limits_{\nu \in \mathbb{N}^n} f_\nu X^\nu$, and ν^0 instead of ν^i, then according to lemma 1.27 (with $\varepsilon = 1$) we can introduce a finite subset $J \subseteq \mathbb{N}^n$ and for $\nu \in J$ some series $\phi_\nu \in \mathcal{A}\{\underline{\rho}^{-1}X\}$ such that $\|\phi_\nu\| < 1$ and

$$f = \sum_{\nu \in J} f_\nu (X^\nu + \phi_\nu) = f_{\nu^0}(X^{\nu^0} + \phi_{\nu^0}) + \left(\sum_{\nu \in J \setminus \{\nu^0\}} f_\nu (X^\nu + \phi_\nu) \right) \tag{6}$$

We then pick a real number $s > 1$ that fulfils the conditions (5), another real number s' such that $s' \in]1, s[\cap \sqrt{|k^*|}$, and consider the following constructible datum of X: $(Y,T) \overset{\varphi}{\dashrightarrow} X$ given by $Y = \mathcal{M}(\mathcal{B})$ where

$$\mathcal{B} = \mathcal{A}\{s'^{-1} t_\nu\}_{\nu \in J \setminus \{\nu^0\}} / (f_\nu - t_\nu f_{\nu^0})$$

(*cf.* the remarks preceding lemma 1.12), and

$$T = \{ y \in Y \bigm| |f_\nu(y)| \leq p |f_{\nu^0}(y)| \text{ and } f_{\nu^0}(y) \neq 0 \}$$

Then $\varphi(T)$ is a neighbourhood of x. Now, since $f_\nu = f_{\nu^0} t_\nu$ for $\nu \in J \setminus \{\nu^0\}$, (6) becomes:

$$\sigma(f) = f_{\nu^0} \left(\sigma(X^{\nu^0}) + \sigma(\phi_{\nu^0}) + \sum_{\nu \in J \setminus \{\nu^0\}} t_\nu (\, \sigma(X^\nu) + \sigma(\phi_\nu)) \right) \tag{7}$$

So if we set

$$g := \sigma(X^{\nu^0}) + \sigma(\phi_{\nu^0}) + \sum_{\nu \in J\setminus\{\nu^0\}} t_\nu(\sigma(X^\nu)) + \sigma(\phi_\nu) \in \mathcal{B}\{\underline{\rho}^{-1}X\} \tag{8}$$

then

$$\sigma(f) = f_{\nu^0} g. \tag{9}$$

We now define d as

$$d := (\max_{\nu\in J} |\nu|_\infty) + 1$$

Hence for $\nu = (\nu_1, \ldots, \nu_n) \in J$

$$\sigma(X^\nu) = (X_1 + X_n^{d^{n-1}})^{\nu_1}(X_2 + X_n^{d^{n-2}})^{\nu_2} \ldots (X_{n-1} + X_n^d)^{\nu_{n-1}}(X_n)^{\nu_n}$$

Since $\nu \leq_{lex} \nu^0$, and by definition of d, for all i and $\nu \in J$, $|\nu_i| < d$, when we develop this expression, we obtain a sum of distinct monomials indexed by decomposition of integers in basis d. Precisely, we obtain exactly the sum of the monomials of the form

$$X_I = (\prod_{l\in I} X_l^{\nu_l})(\prod_{l\in\{1\ldots n-1\}\setminus I} X_n^{\nu_l d^{n-l}})X_n^{\nu_n} = (\prod_{l\in I\cup\{n\}} X_l^{\nu_l})X_n^{\sum_{l\in\{1\ldots n-1\}\setminus I}\nu_l d^{n-l}}$$

where I is any subset of $\{1\ldots n-1\}$. One checks that by definition of $\underline{\rho}$, $\|X_I\|_{\underline{\rho}} = s^{\left(\sum\limits_{k=1}^{n}\nu_k d^{n-k}\right)}$. As a consequence, it is independent of I and depends only on ν. Moreover, if we count in base d, since ν^0 is maximal in J according to the lexicographic order, for all $\nu \in J \setminus \{\nu^0\}$

$$\sum_{k=1}^{n} \nu_k d^{n-k} + 1 \leq \sum_{k=1}^{n} \nu_k^0 d^{n-k}$$

Hence the monomial

$$X_n^{\sum\limits_{k=1}^{n}\nu_k^0 d^{n-k}}$$

which appears in $\sigma(X^{\nu^0})$ fulfils for all $\nu \in J \setminus \{\nu^0\}$:

$$\|\sigma(X^{\nu^0})\|_{\underline{\rho}} = \|X_n^{\sum\limits_{k=1}^{n}\nu_k^0 d^{n-k}}\|_{\underline{\rho}} = s^{\sum\limits_{k=1}^{n}\nu_k^0 d^{n-k}} \geq s^{1+\sum\limits_{k=1}^{n}\nu_k d^{n-k})} \geq s\|\sigma(X^\nu)\|_{\underline{\rho}} > s'\|\sigma(X^\nu)\|_{\underline{\rho}} \geq \|t_\nu\sigma(X^\nu)\|_{\underline{\rho}}$$

(according to the definition of Y).
Now remind that $\sigma(f) = f_{\nu^0}g$ (9), where we set in (8)

$$g = \sigma(X^{\nu^0}) + \sigma(\phi_{\nu^0}) + \sum_{\nu\in J\setminus\{\nu^0\}} t_\nu(\sigma(X^\nu)) + \sigma(\phi_\nu) \in \mathcal{B}\{\underline{\rho}^{-1}X\}.$$

But $\sigma(X^{\nu^0}) = X_n^{\sum\limits_{k=1}^{n}\nu_k^0 d^{n-k}} + \ldots$ where the remaining terms appearing in these $\ldots$ are of degree strictly less than $\sum\limits_{k=1}^{n}\nu_k^0 d^{n-k}$ in X_n. And all have the same norm in $\mathcal{B}\{\underline{\rho}^{-1}X\}$, namely $s^{\sum_{k=1}^{n}\nu_k^0 d^{n-k}}$.
By hypothesis, in $\mathcal{A}\{\underline{\rho}^{-1}X\}$, $\|\phi_\nu\| < 1$ for all $\nu \in J$. Since $\underline{\rho} < \underline{r}$, and $\mathcal{A} \to \mathcal{B}$ is contractive, this inequality remains true in $\mathcal{B}\{\underline{\rho}^{-1}X\}$.
In conclusion $\|\sigma(X^{\nu^0})\|_{\underline{\rho}} > \|\sigma(\phi_\nu)\|_{\underline{\rho}}$ for $\nu \in J$, and $\|\sigma(X^{\nu^0})\|_{\underline{\rho}} > \|t_\nu\sigma(X^\nu)\|_{\underline{\rho}}$ for $\nu \in J \setminus \{\nu^0\}$. So g is X_n distinguished of order $\sum\limits_{k=1}^{n}\nu_k^0 d^{n-k}$ in $\mathcal{B}\{\underline{\rho}^{-1}X\}$.

Now remind that in fact $f = f_i$, and $\nu^0 = \nu^i$. Hence we have shown that $\sigma(f_i) = f^i_{\nu^i} g_i$ with g_i X_n-distinguished, and if we set $\beta_i = f^i_{\nu^i}$ we have the result we wanted to prove, except that we made it for a specific i. But if we take a d big enough that works for all i, and $s > 1$ close enough to 1 that also works for all i, and if, using corollary 1.10, we define (Y, T) as the fibered product of all the (Y_i, T_i) we are done (there are things to check, for instance that X_n-distinguished functions remain X_n-distinguished after this fibered product, which is remark 1.23).

□

Theorem 1.31. *Let $S \subset X$. Then S is an overconvergent subanalytic subset if and only if it is an overconvergent constructible subset.*

Proof. Let's first suppose that S is an overconvergent subanalytic subset. Then by definition there exist $r > 1$, T a globally semianalytic subset of $X \times \mathbb{B}_{\underline{r}}$ such that $S = \pi\left(T \cap (X \times \mathbb{B}^n)\right)$. We then show the result by induction on n.
If $n = 0$, there is nothing to prove since by definition, X is then a globally semianalytic subset of X, in particular it is an overconvergent constructible subset.
Let then $n > 0$ be given and assume that the theorem holds for integers $< n$. In order to prove the theorem, we can actually assume that T is a basic semianalytic subset, i.e. that there are $2m$ functions $f_1, \ldots, f_m, g_1, \ldots, g_m \in \mathcal{A}\{\underline{r}^{-1}X\}$ and $\Diamond_j \in \{\leq, <\}$ for $j = 1 \ldots m$ such that

$$T = \{x \in X \times \mathbb{B}^n_r \bigm| |f_j(x)| \Diamond_j |g_j(x)| \ j = 1 \ldots m\}$$

We can make this assumption because any globally semianalytic subset of $X \times \mathbb{B}^n_r$ is a finite union of basic semianalytic subsets. Now, according to proposition 1.28 we can find a constructible covering $(X_i, S_i) \overset{\varphi_i}{\dashrightarrow} (X, S)$ where $X_i = \mathcal{M}(\mathcal{B}_i)$ such that on each S_i, and for all j, $f_j = a^i_j F^i_j$ and $g_j = b^i_j G^i_j$, where $a^i_j, b^i_j \in \mathcal{B}_i$, $F^i_j, G^i_j \in \mathcal{B}_i\{\underline{r}^{-1}X\}$, and the coefficients of F^i_j (and of G^i_j) generate the unit ideal. Then on each X_i we apply the proposition 1.30 to the functions F^i_j and G^i_j. By compactness of each X_i, we can find for each i a constructible covering of X_i such that the conclusions of proposition 1.30 are true. Now if we compose this succession of constructible covers (of X first, and then of the X_i's) we can assert that:
there exists a constructible covering $(X_i, S_i) \overset{\varphi_i}{\dashrightarrow} X$, $i = 1 \ldots N$, with $X_i = \mathcal{M}(\mathcal{B}_i)$, and for each $i = 1 \ldots N$ some polyradii $\underline{\rho_i}$ such that $\underline{1} < \underline{\rho_i} \leq \underline{r}$, and some Weierstrass automorphisms σ_i of $X_i \times \mathbb{B}_{\underline{\rho_i}}$ such that if we consider the following diagram

$$\begin{tikzcd}
 & (X_i, S_i) \arrow[r, dashed, "\varphi_i"] & X \\
X_i \times \mathbb{B}_{\underline{\rho_i}} \arrow[ur, "\pi_i"] \arrow[r, "\sigma_i"] & X_i \times \mathbb{B}_{\underline{\rho_i}} \arrow[u, "\pi_i"] \arrow[r, dashed, "\varphi'_i"] & X \times \mathbb{B}_{\underline{\rho_i}} \arrow[u, "\pi"]
\end{tikzcd}$$

and if we define

$$T_i := \varphi_i'^{-1}(T) \subseteq X_i \times \mathbb{B}_{\underline{\rho_i}}$$

then

$$\pi(T) = \cup_{i=1}^N \varphi_i(\pi_i(T_i) \cap S_i) \tag{10}$$

because $(X_i, S_i) \overset{\varphi_i}{\dashrightarrow} X$, $i = 1 \ldots N$ is a constructible cover. Hence, using proposition 1.11(2) we can now restrict to prove that $\pi_i(T_i)$ is an overconvergent constructible subset.
We set $\alpha_i = \varphi'_i \circ \sigma_i$. Then $\alpha^*_i(f_j)_{|\pi_i^{-1}(S_i)} = a^i_j F^i_{j\,|\pi_i^{-1}(S_i)}$ (resp. $\alpha^*_i(g_j)_{|\pi_i^{-1}(S_i)} = b^i_j G^i_{j\,|\pi_i^{-1}(S_i)}$) with a^i_j (resp. b^i_j) $\in \mathcal{B}_i$ and F^i_j (resp. G^i_j) a X_n-distinguished element of $\mathcal{B}_i\{\underline{\rho_i}^{-1}X\}$. Then according to Weierstrass preparation (proposition 1.26) applied to F^i_j and G^i_j, we can write $F^i_j = e^i_j w^i_j$ where e^i_j is a multiplicative unit of $\mathcal{B}_i\{\underline{r}^{-1}X\}$, and w^i_j a unitary polynomial of $\mathcal{B}_i\{\rho_1^{-1}X_1, \ldots, (\rho_{n-1})^{-1}X_{n-1}\}[X_n]$. Thus, $P^i_j := a^i_j w^i_j$ is a polynomial in the sense that it belongs to $\mathcal{A}\{r_1^{-1}X_1, \ldots, (r_{n-1})^{-1}X_{n-1}\}[X_n]$, and on $\pi_i^{-1}(S_i)$, $f^i_j = e^i_j P^i_j$ (resp $g^i_j = E^i_j Q^i_j$) where e^i_j and E^i_j are multiplicative units and P^i_j, Q^i_j polynomials. Then (*cf.* remark 1.20), for all x in X_i, $|e^i_j(x)| = \|e^i_j\|$ (resp. $|E^i_j(x)| = \|E^i_j\|$) and moreover $\|e^i_j\|$ and $\|E^i_j\|$ belong to $\sqrt{|k^\times|}$. So

$$T_i = \{x \in X_i \times \mathbb{B}_{\underline{r}} \bigm| |f^i_j(x)| \Diamond_j |g^i_j(x)|,\ j = 1 \ldots m\} = \{x \in X_i \times \mathbb{B}_{\underline{r}} \bigm| \|e^i_j\| |P^i_j(x)| \diamond_j \|E^i_j\| |Q^i_j(x)|,\ j = 1 \ldots m\}$$

Let's then consider the projection on the $n-1$ coordinates of $\mathbb{B}_{\underline{r}}$:

$$X_i \times \mathbb{B}_{\underline{r}} \xrightarrow{\pi'_i} X_i \times \mathbb{B}_{(r_1, \ldots, r_{n-1})} \xrightarrow{\pi''_i} X_i$$

According to [8, 2.5] (where one of the main tool is quantifier elimination in the theory of algebraically closed valued fields), $\pi'_i(T_i)$ is a globally semianalytic subset of $X_i \times \mathbb{B}_{((r_1, \ldots r_{n-1})}$ so by induction hypothesis on n we can conclude that $\pi_i(T_i) = \pi''_i((\pi'_i(T_i))$ is an overconvergent constructible subset of X_i, and with (10) this ends this part of the proof.

Conversely, let's show that if $S \subset X$ is an overconvergent constructible subset, it is also overconvergent subanalytic.

Let $(Y,T) \xrightarrow{\varphi} (X,S)$ be an elementary constructible datum associated with $f, g \in \mathcal{A}$, $r > s > 0$ and $R \subset Y$ a globally semianalytic subset. So that $T = \{y \in R \mid |f(y)| \leq s|g(y)| \neq 0\}$. Moreover, let U be an overconvergent subanalytic subset of Y such that $U \subseteq T$, defined with $(r_1, \ldots, r_n) = \underline{r} > \underline{s} > 0$, $\underline{s} \in \sqrt{|k^\times|}^n$, W a globally semianalytic subset of $Y \times \mathbb{B}_{\underline{r}}$ and $U = \pi(W \cap (X \times \mathbb{B}_{\underline{s}})$. Then let's show that $\varphi(U)$ is an overconvergent subanalytic subset of X.
Indeed by definition, $Y = \mathcal{M}(\mathcal{B})$ where $\mathcal{B} = \mathcal{A}\{r^{-1}t\}/(f-tg)$. Hence $Y \times \mathbb{B}_{\underline{r}}$ can naturally be identified with the Zariski closed subset $V(f-tg)$ of $X \times \mathbb{B}_{(r,r_1,\ldots,r_n)} = X \times \mathcal{M}(k\{r^{-1}t, r_1^{-1}t_1, \ldots, r_n^{-1}t_n\})$. We then obtain the following commutative diagram:

$$\begin{tikzcd}
Y \arrow[r, hook] & X \times \mathbb{B}_r \arrow[r] & X \\
Y \times \mathbb{B}_{\underline{r}} \arrow[u] \arrow[r, hook, "\alpha"] & X \times \mathbb{B}_{(r,\underline{r})} \arrow[u] \arrow[ru, "\pi"] &
\end{tikzcd}$$

where the arrows $\hookrightarrow$ are embeddings of Zariski closed subsets. Then $W' := \alpha(W)$ is a globally semianalytic subset of $X \times \mathbb{B}_{(r,\underline{r})}$ (see lemma 1.41), it then follows that $\varphi(U) = \pi(W' \cap X \times (\mathbb{B}_{(s,\underline{s})})$ is an overconvergent subanalytic subset of X (see remark 1.15).
From this remark, one proves easily by induction on the complexity of a constructible datum $(Y,T) \overset{\varphi}{\dashrightarrow} X$ that if U is an overconvergent subanalytic subset of Y, $\varphi(U \cap T)$ is an overconvergent subanalytic subset of X. In particular, if U is a globally semianalytic subset of Y, then $\varphi(U \cap T)$ is overconvergent subanalytic in X, which is what we wanted to prove. ☐

Hence thanks to this theorem we can use some obvious properties of overconvergent subanalytic (resp. constructible) subsets to prove less obvious results about overconvergent constructible (resp. subanalytic) subsets. For instance we can obtain the non-trivial result concerning overconvergent subanalytic subsets:

Proposition 1.32. *Let X be an affinoid space, then the class of overconvergent subanalytic subsets of X is stable under finite boolean combination*[3]*.*

Proof. This was proven for overconvergent constructible subsets in proposition1.11. ☐

In the same way, we obtain here a non-obvious stability property of overconvergent constructible subsets:

Corollary 1.33. *Let $\underline{r} \in \mathbb{R}^n$ be a polyradius such that $\underline{r} > \underline{1}$, and $S \subseteq X \times \mathbb{B}_{\underline{r}}$ be an overconvergent subanalytic (or constructible) subset of $X \times \mathbb{B}_{\underline{r}}$. Then $\pi\left(S \cap (X \times \mathbb{B}^n)\right)$ is an overconvergent subanalytic (or constructible) subset of X.*

Proof. If S is an overconvergent subanalytic subset of $X \times \mathbb{B}_{\underline{r}}$, by definition, there exists $s > 1$, an integer m and T a globally semianalytic subset of $X \times \mathbb{B}_{\underline{r}} \times \mathbb{B}_s^m$ such that $S = \pi_2(T \cap ((X \times \mathbb{B}_{\underline{r}}) \times \mathbb{B}^m))$ where $\pi_2 : (X \times \mathbb{B}_{\underline{r}}) \times \mathbb{B}_s^m \to X \times \mathbb{B}_{\underline{r}}$ is the natural projection. Hence $\pi(S \cap (X \times \mathbb{B}^n)) = \pi_2(T \cap ((X \times \mathbb{B}^n) \times \mathbb{B}^m)) = \pi_2(T \cap (X \times \mathbb{B}^{n+m})$ where $\pi_2 : X \times \mathbb{B}_{\underline{r}} \times Bs^m \to X$ is the natural projection (so $\pi_2 = \pi \circ \pi_1$). Hence S is an overconvergent subanalytic subset of X. ☐

1.5. From a global to a local definition.

From now on, all k-analytic spaces will be good k-analytic spaces (these are the k-analytic spaces such that any point has an affinoid neighbourhood, see [2, 1.2.16]).

Definition 1.34. Let X be a k-analytic space. A wide covering of X is a covering $\{X_i\}$ such that the $X_i's$ are affinoid domains of X and $\{\overset{\circ}{X_i}\}$ is also a covering of X.

Proposition 1.35. *Let X be an affinoid space, and S a subset of X. The following assertions are equivalent:*

1. *S is an overconvergent subanalytic subset of X.*
2. *For all wide covering $\{X_i\}$ of X, $X_i \cap S$ is an overconvergent subanalytic subset of X_i.*
3. *There exists a wide covering $\{X_i\}$ of X such that $X_i \cap S$ is overconvergent subanalytic in X_i for all i.*

---

[3] In fact, the only non-trivial result is that overconvergent subanalytic subsets are stable under taking complementary.

(4) *For all $x \in X$ there exists an affinoid neighbourhood V of x such that $V \cap S$ is overconvergent subanalytic in V.*

This last equivalence implies that the class of overconvergent subanalytic subsets is local in the sense of definition 0.1.

Proof. (1) $\Rightarrow$ (2) is a consequence of lemma 1.14.
(2) $\Rightarrow$ (3) is clear.
(3) $\Leftrightarrow$ (4) follows from the compactness of X.
(4) $\Rightarrow$ (1) Let $x \in X$ and V an affinoid neighbourhood of x such that $V \cap S$ is overconvergent subanalytic in V. Since rational domains are a basis of neighbourhood of x, we can assume that V is a a rational domain. Then, $V \cap S$ is still an overconvergent subanalytic subset of V. According to [1, 2.5.15], there exists a presentation of this rational domain :

$$V = \{z \in X \mid |f_j(z)| \leq r_j |g(z)| \ j = 1 \dots n\}$$

and some real numbers $p_1 \dots p_n$ such that $p_j \in \sqrt{k^*}$, $0 < p_j < r_j$ for all j and $x \in \mathring{U_x}$ where $U_x = \{z \in X \mid |f_j(z)| \leq p_j |g(z)| \ j = 1 \dots n\}$. It is then easy to check that $(V, U_x) \overset{\iota}{\dashrightarrow} X$ is a constructible datum (cf. lemma 1.12 and the remark preceding it). Since $V \cap S$ is also overconvergent constructible in V, $U_x \cap S$ is overconvergent constructible (hence also overconvergent subanalytic) in X according to proposition 1.11(2). By compactness, we can extract a finite covering from the U_x's so S is overconvergent subanalytic in X. ☐

Definition 1.36. Let X be a k-analytic space. A subset $S \subset X$ is called overconvergent subanalytic (or overconvergent constructible) if for all $x \in X$ there exists an affinoid neighbourhood V of x such that $V \cap S$ is overconvergent subanalytic in V.

According to the last proposition, when X is an affinoid space, this definition is compatible with the previous one (definition 1.13).

Remind the definition of a locally semianalytic subset that we have given in the introduction:

Definition 1.37. Let X be a good k-analytic space. A subset S of X is locally semianalytic if for all $x \in X$ there exists an affinoid neighbourhood V of x such that $S \cap V$ is globally semianalytic in V .

Corollary 1.38. *Let X be a Hausdorff k-analytic space. The class of locally semianalytic subsets of X is contained in the class of overconvergent constructible subsets of X.*

Corollary 1.39. *Let X be an affinoid space, then $S \subset X$ is an overconvergent subanalytic (or constructible) subset of X if and only if there exist $r > 1$, an integer n, and $T \subseteq X \times \mathbb{B}^n_r$ a locally semianalytic subset, such that $S = \pi(T \cap (X \times \mathbb{B}^n))$.*

Proof. The first implication is true because a globally semianalytic subset of $X \times \mathbb{B}^n_r$ is in particular a locally semianalytic subset of $X \times \mathbb{B}^n_r$.
Conversely, if $S = \pi(T \cap (X \times \mathbb{B}^n))$ where T is a locally semianalytic subset of $X \times \mathbb{B}^n_r$, then according to corollary 1.38, T is overconvergent subanalytic in $X \times \mathbb{B}^n_r$, so according to corollary 1.33, $\pi(T \cap (X \times \mathbb{B}^n))$ is also overconvergent subanalytic. ☐

Lemma 1.40. *Let $\varphi : X \to Y$ be a morphism of k-analytic spaces, and $S \subseteq Y$ be a locally semianalytic subset. Then $\varphi^{-1}(S)$ is locally semianalytic.*

Proof. Let $x \in X$, $y = \varphi(x)$. There exists an affinoid neighbourhood V of Y such that $V \cap S$ is globally semianalytic in V. Let W be an affinoid neighbourhood of x in $\varphi^{-1}(V)$. Then $W \cap \varphi^{-1}(S)$ is globally semianalytic in W. ☐

Lemma 1.41. *Let $\varphi : X \to Y$ be a closed immersion.*

(1) *If φ is a morphism of affinoid spaces, and S is a globally semianalytic subset of X, then $\varphi(S)$ is a globally semianalytic subset of Y.*
(2) *Let S be an overconvergent constructible subset of X, then $\varphi(S)$ is an overconvergent constructible subset of Y.*

Proof.

(1) Write $Y = \mathcal{M}(\mathcal{A})$ and $X = \mathcal{M}(\mathcal{A}/\mathcal{I})$ where $\mathcal{I} = (a_1, \dots, a_m)$ is an ideal of $\mathcal{A}$. Then, if $S = \{x \in X \mid |f_i(x)|\Diamond_i|g_i(x)|,\ i = 1 \dots n\}$ with $f_i, g_i \in \mathcal{A}/\mathcal{I}$, we can find functions $F_i, G_i \in \mathcal{A}$ such that $\overline{F_i} = f_i$ and $\overline{G_i} = g_i$. In that case one checks that, $\varphi(S) = \{y \in Y \mid |F_i(y)|\Diamond_i|G_i(y)|,\ i = 1 \dots n\} \cap \{y \in Y \mid a_i(y) = 0,\ i = 1 \dots m\}$, which is indeed globally semianalytic.

(2) Since the problem is local on Y, we can assume that Y is affinoid, and hence, X is also affinoid. Using the equivalence of theorem 1.31 it is equivalent to show that if S is an overconvergent subanalytic subset of X, so is $\varphi(S)$. By definition, this means that there exists $T \subseteq X \times \mathbb{B}^n_r$ for some $r > 1$ such that $S = \pi(T \cap (X \times \mathbb{B}^n)$. We then consider the following cartesian diagram:

$$\begin{tikzcd}
X \times \mathbb{B}^n_r \arrow[r, "\varphi'"] \arrow[d, "\pi'"] & Y \times \mathbb{B}^n_r \arrow[d, "\pi"] \\
X \arrow[r, "\varphi"] & Y
\end{tikzcd}$$

But φ' is also a closed immersion, so according to (1), $T' = \varphi'(T)$ is a globally semianalytic subset of $Y \times \mathbb{B}^n_r$. Then one checks that

$$\pi\left(T' \cap (Y \times \mathbb{B}^n)\right) = \pi\left(\varphi'(T) \cap (Y \times \mathbb{B}^n)\right) = \pi\left(\varphi'(T \cap (X \times \mathbb{B}^n)\right) = \varphi(\pi'(T \cap (X \times \mathbb{B}^n)) = \varphi(S).$$

□

Theorem 1.42. *Let $\varphi : Y \to X$ be a morphism of affinoid spaces, and U an affinoid domain of Y such that $U \subseteq Int(Y/X)$. If S is an overconvergent constructible subset of Y then $\varphi(U \cap S)$ is an overconvergent constructible subset of X.*

Proof. According to [1, Prop 2.5.9] there exist $\underline{r} > \underline{s} > 0$ and $\mathcal{A}\{\underline{r}^{-1}t\} \to \mathcal{B}$ an admissible epimorphism which hence identifies Y with a Zariski closed subset of $X \times \mathbb{B}_{\underline{r}}$, such that under this identification, $U \subseteq X \times \mathbb{B}_{\underline{s}}$. We can assume that $\underline{s} \in \sqrt{|k^\times|}^n$. If we note $\Gamma(\varphi)$ the graph of φ, this induces a Zariski closed embedding of $Y \simeq \Gamma(\varphi) \xrightarrow{i} X \times \mathbb{B}_{\underline{r}}$. Now since S is an overconvergent constructible subset of Y, according to lemma 1.41, $i(S)$ is an overconvergent constructible subset of $X \times \mathbb{B}_{\underline{r}})$. Finally, U is a globally semianalytic subset of Y (because of Gerritzen-Grauert theorem), so $i(U)$ is also globally semianalytic in $X \times \mathbb{B}_{\underline{r}}$, and by assumption, $i(U) \subseteq X \times \mathbb{B}_{\underline{s}}$, so $i(U \cap S) \subseteq X \times \mathbb{B}_{\underline{r}}$ is an overconvergent constructible subset of $X \times \mathbb{B}_{\underline{r}}$, and according to corollary 1.33, $\pi(i(U \cap S))$ is an overconvergent constructible subset of X. But this set is precisely $\varphi(U \cap S)$. □

Proposition 1.43. *Let $\varphi : Y \to X$ be a morphism of k-analytic spaces, S an overconvergent constructible subset of Y such that $\overline{|S|} \to |X|$ is proper and $\overline{|S|} \subseteq Int(Y/X)$. Then $\varphi(S)$ is an overconvergent constructible subset of X.*

Proof. If X' is an affinoid domain of X and if we consider the cartesian diagram :

$$\begin{tikzcd}
S \subseteq Y \arrow[r, "\varphi"] & X \\
S' \subseteq Y' \arrow[r, "\varphi'"] \arrow[u, "\psi'"] & X' \arrow[u, "\psi"']
\end{tikzcd}$$

then $\psi'^{-1}(\overline{S})$ is closed in Y' and contains $\psi'^{-1}(S) = S'$ so $S' \subseteq \overline{S'} \subseteq \psi'^{-1}(\overline{S})$, and since properness is stable under base change, $\psi'^{-1}(\overline{S}) \to |X'|$ is proper, and since $\overline{S'}$ is closed, $\overline{S'} \to |X'|$ is proper. Moreover, $\psi'^{-1}(\mathrm{Int}(Y/X)) \subseteq \mathrm{Int}(Y'/X')$ ([1, 3.1.3 (iii)]) so $\overline{S'} \subseteq \psi'^{-1}(\overline{S}) \subseteq \mathrm{Int}(Y'/X')$. So S' and φ' fulfil the hypotheses of the proposition. Hence, since the property we want to check is local on X, we can assume that X is an affinoid space, hence that $\overline{|S|}$ is compact.

Now for every $y \in \overline{|S|}$ we can find an affinoid neighbourhood U such that $U \subseteq \mathrm{Int}(Y/X)$, because $\mathrm{Int}(Y/X)$ is open. Then, $\varphi(U \cap S)$ is an overconvergent constructible subset of X according to theorem 1.42. Since $\overline{|S|}$ is compact we can extract from this a finite covering of $\overline{|S|}$, which finishes to prove that $\varphi(S)$ is overconvergent constructible. □

Corollary 1.44. *Let* $\varphi : Y \to X$ *be a proper morphism of good* k*-analytic spaces. Remind ([1, p.50]) that it means that* $|\varphi| : |Y| \to |X|$ *is proper and that* $\partial(Y/X) = \emptyset$*. Let* S *be an overconvergent constructible subset of* Y*. Then* $\varphi(S)$ *is an overconvergent constructible subset of* X*.*

Definition 1.45. A morphism $\varphi : Y \to X$ of good k-analytic spaces is locally extendible without boundary if, for all $y \in Y$, there exists an affinoid neighbourhood U of y, Y' an affinoid space that contains U as an affinoid domain, and $\psi : Y' \to X$ that extends $\varphi_{|U}$, such that $U \subseteq \mathrm{Int}(Y'/X)$.

Remark that using again [1, 3.1.3 (iii)], this property is stable under base change.

Proposition 1.46. *Let* $\varphi : Y \to X$ *be a compact morphism (i.e.* $|\varphi| : |Y| \to |X|$ *is proper [1, p.50]) which is locally extendible without boundary. Then* $\varphi(Y)$ *is an overconvergent constructible subset of* X*.*

Proof. We can assume that X is an affinoid space, so Y is compact. Then for all $y \in Y$ we can find an affinoid neighbourhood U of y and Y' an affinoid space that contains U, and $\psi : Y' \to X$ that extends $\varphi_{|U}$, such that $U \subseteq \mathrm{Int}(Y'/X)$. Then, according to theorem 1.42, $\varphi(U)$ is an overconvergent constructible subset of X (take $S = Y'$). Hence by compactness of Y, $\varphi(Y)$ is overconvergent constructible. ☐

2. Study of various classes

2.1. **Many families.** In this section $X = \mathcal{M}(\mathcal{A})$ will be an affinoid space.
The aim of this section is to first recall the definitions of the various classes of *rigid/locally/strongly/D-semianalytic/subanalytic* subsets of X that are defined in [19].
We now give the following definitions. A subset $S \subseteq X$ is called :

(a) Globally semianalytic if it is a boolean combination of subsets of the form $\{x \in X \mid |f(x)| \leq |g(x)|\}$, with $f, g \in \mathcal{A}$.
(b) Locally semianalytic, if for all x in X there exists an affinoid neighbourhood V of x such that $S \cap V$ is globally semianalytic in V.
(c) Rigid semianalytic if there is a finite affinoid covering $\{X_i\}_{i=1}^n$ such that $S \cap X_i$ is globally semianalytic in X_i for all i.
(d) Overconvergent subanalytic has been defined in definition 1.6. As we proved in the previous section, this corresponds also to overconvergent constructible subsets. In [19]. Moreover, our definition of overconvergent subanalytic subset is the same as the definition of globally strongly subanalytic of [19, 1.3.8.1]. In [19] it is proven that this is equivalent to the class of globally strongly **D**-semianalytic subsets [19, 1.3.2].
(e) Rigid overconvergent constructible if there exists a finite affinoid covering $\{X_i\}$ of X such that $S \cap X_i$ is overconvergent constructible in X_i for all i. This corresponds to the notion of *strongly* **D** *-semianalytic* subset in [19, 1.3.7.1].
(f) Strongly subanalytic if there exists an integer n, $r > 1$, a subset $T \subseteq X \times \mathbb{B}_r^n$ which is rigid semianalytic, such that $S = \pi(T \cap (X \times \mathbb{B}^n)$. This is definition [19, 1.3.8.1], and we'll give an equivalent definition in proposition 2.8.
(g) Locally strongly subanalytic if there exists a finite affinoid covering $\{X_i\}$ of X such that $S \cap X_i$ is strongly subanalytic in X_i for all i. This is definition [19, 1.3.8.2].

In [19] it is stated that (d),(e),(f) and (g) are equivalent (equivalence of $(e), (f), (g)$ is stated in [19, Prop 4.2] , and the equivalence of (d) and (f) is stated in [19, Th 5.2]). These results rest on [19, lemma 4.1] which is false, and we'll show indeed that $(d), (e)$ and (f) correspond in general to three different classes. More precisely the aim of this section is to show that these classes satisfy the following relations:

FIGURE 1. The hierarchy



In this figure, Class A $\supsetneq$ Class B, means that the class A *properly* contains the class B, and Class A $\not\supseteq$ Class B means that the Class A does not contain the class B.

In this diagram, all the inclusions are clear from the definitions (except that the class of overconvergent constructible subsets contains the class of locally semianalytic subsets, which is corollary 1.35). In comparison with what was stated in [19], the most striking inequality is probably $\not\subseteq^5$ which asserts that rigid semianalytic subsets are not necessarily overconvergent subanalytic subsets whereas according to [19, Th 5.2], they should be overconvergent subanalytic. In other words, when you project overconvergent globally semianalytic subsets, you obtain a class which is not G-local (but however local for the Berkovich topology).

In this section we'll show that the inclusions (1)-(8) in figure 1 are all proper (in the next section we'll explain that if X is regular of dimension 2, overconvergent subanalytic subsets correspond to locally semianalytic subsets). We don't know if the inclusion on the left

$$\textit{locally strongly subanalytic} \supseteq \textit{strongly subanalytic}$$

is proper.

2.2. Rigid semianalytic subsets are not necessarily overconvergent constructible.

Here we prove the inequality $\not\subseteq^5$.

Lemma 2.1. *Let $\eta \in X$ such that $\mathcal{O}_{X,\eta}$ is a field, $S \subset X$ a globally semianalytic subset. If $\eta \in \overline{S}$, then $\overset{\circ}{S}$ is non empty.*

Proof. Since

$$\bigcup_{i=1}^{n} \overline{S_i} = \overline{\bigcup_{i=1}^{n} S_i}$$

we can assume that S is a basic semianalytic subset, i.e is of the form:

$$S = \left(\bigcap_{i=1}^{m} \{x \in X \mid |f_i(x)| \leq |g_i(x)|\} \right) \bigcap \left(\bigcap_{j=1}^{n} \{x \in X \mid |F_j(x)| < |G_j(x)|\} \right)$$

We use the following decomposition

$$\{x \in X \mid |f_i(x)| \leq |g_i(x)|\} = \{x \in X \mid f_i(x) = 0\} \cup \{x \in X \mid f_i(x) \neq 0 \text{ and } |f_i(x)| \leq |g_i(x)|\}$$

and using again that the adherence is stable under finite union, we can assume that $\eta \in \overline{S}$ and that S is of the form:

$$S = \bigcap_{i=1}^{l} \{x \in X \mid h_i(x) = 0\} \bigcap_{j=1}^{m} \{x \in X \mid f_j(x) \neq 0 \text{ and } |f_j(x)| \leq |g_j(x)|\} \bigcap_{k=1}^{n} \{x \in X \mid |F_k(x)| < |G_k(x)|\}$$

Since the subsets $\{x \in X \mid h_i(x) = 0\}$ are closed, and contain S and $\eta \in \overline{S}$, $h_i(\eta) = 0$.
Since $\mathcal{O}_{X,\eta}$ is a field we can find an affinoid neighbourhood V of η such that $h_{i|V} = 0$ for all i. Hence $V \cap S \neq \emptyset$ (because $\eta \in \overline{S}$) and we can remove the $h_i's$, and assume that

$$V \cap S = \bigcap_{j=1}^{m} \{x \in V \mid f_j(x) \neq 0 \text{ and } |f_j(x)| \leq |g_j(x)|\} \bigcap_{k=1}^{n} \{x \in V \mid |F_k(x)| < |G_k(x)|\}$$

This defines a strictly k-analytic domain of X, which is non empty, so its interior is also non empty, for instance its interior contains some rigid points. $\square$

Lemma 2.2. *Let $\eta \in X$ and assume that $\mathcal{O}_{X,\eta}$ is a field. Let $(Y,T) \xrightarrow{\varphi} (X,S)$ be an elementary constructible datum with $Y = \mathcal{M}(\mathcal{A}\{r^{-1}t\}/(f-tg))$ where $T = \varphi^{-1}(S) \cap \{y \in R \mid g(y) \neq 0$ and $|f(y)| \leq s|g(y)|\}$ with $0 < s < r$, $s \in \sqrt{|k^\times|}$ and R a globally semianalytic subset of Y. Let's assume that $\eta \in \overline{\varphi(T)}$. Then*

(a) $g(\eta) \neq 0$

(b) $|f(\eta)| \leq s|g(\eta)|$

(c) There exists a neighbourhood U of η such that $\varphi^{-1}(U) \xrightarrow{\varphi_{|\varphi^{-1}(U)}} U$ is an isomorphism. If we note η' the only point of $\varphi^{-1}(U)$ such that $\varphi(\eta') = \eta$, then $\eta' \in \overline{T}$ and $\mathcal{O}_{Y,\eta'}$ is a field.

Proof.

(a) If we had $g(\eta) = 0$, since $\mathcal{O}_{X,\eta}$ is a field, there would exist an affinoid neighbourhood of η, V, such that $g_{|V} = 0$. Since for $p \in T$, $g(\varphi(p)) \neq 0$ we should have $\varphi(T) \cap V = \emptyset$ which is impossible since $\eta \in \overline{\varphi(T)}$.

(b) The subset $\{x \in X \mid |f(x)| \leq s|g(x)|\}$ is a closed subset of X which contains $\varphi(T)$, hence by assumption also η.

(c) If we note $U = \{y \in Y \mid g(y) \neq 0\}$, $\varphi_{|U}$ identifies through an isomorphism U with $\varphi(U) = \{x \in X \mid g(x) \neq 0$ and $|f(x)| \leq r|g(x)|\}$ which is an analytic domain of X, and a neighbourhood of η according to the two preceding points. So $\eta \in \varphi(U)$, let's say $\eta = \varphi(\eta')$ with $\eta' \in U$. Now, $\mathcal{O}_{Y,\eta'} \simeq \mathcal{O}_{X,\eta}$ is a field and $\eta' \in \overline{T}$.

$\square$

Corollary 2.3. *Let $\eta \in X$ such that $\mathcal{O}_{X,\eta}$ is a field, and let U be an overconvergent constructible subset of X. If $\eta \in \overline{U}$, then $\overset{\circ}{U} \neq \emptyset$.*

Proof. Using similar arguments as in the beginning of lemma 2.1, we can assume that $U = \varphi(T)$ where $(Y,T) \overset{\varphi}{\dashrightarrow} X$ is a constructible datum. Hence T is a globally semianalytic subset of Y. A repeated use of lemma 2.2 allows us to say that there exists an open neighbourhood U of η, such that $\varphi^{-1}(U) \xrightarrow{\varphi_{|\varphi^{-1}(U)}} U$ is an isomorphism. Thanks to lemma 2.2 again, we can introduce η', the only point of $\varphi^{-1}(U)$ such that $\varphi(\eta') = \eta$, assert that $\mathcal{O}_{Y,\eta'}$ is a field and that $\eta' \in \overline{T}$. Now if we consider V a strictly affinoid neighbourhood of η' contained in $\varphi^{-1}(U)$, it is true that $\eta' \in \overline{T \cap V}$ (the adherence is here taken in V). Now, $T \cap V$ is a globally semianalytic subset of V so according to lemma 2.1, $T \cap V$ has non empty interior in V. We can then deduce that T has non empty interior in X whence $\varphi(T)$ has also non-empty interior. $\square$

Proposition 2.4. *The subset S of example 0.5 is rigid semianalytic but not overconvergent subanalytic. As a consequence, the class of overconvergent subanalytic subsets is not G-local.*

Proof. Remind that in this example $X = \mathbb{B}^2 = \mathcal{M}(k\{T_1,T_2\})$ is the closed bidisc, $0 < r < 1$ with $r \in |k^*|$, say $r = |\varepsilon|$, and $f \in k\{r^{-1}u\}$ whose radius of convergence is exactly r, and $\|f\| < 1$. We then defined

$$S = \{x \in X \mid |T_1(x)| < r \text{ and } T_2(x) = f(T_1(x))\}$$

In more concrete terms, S is the set of points of the *curve* whose equation is $T_2 = f(T_1)$, restricted to the subset $\{|T_1| < r\}$. Let's consider

$$\begin{array}{ccc} \mathbb{B} & \xrightarrow{\psi} & X \\ u & \mapsto & (\varepsilon u, f(\varepsilon u)) \end{array}$$

and let's note $\eta = \psi(g)$ where g is the Gauss point of $\mathbb{B}$. Then $S \subseteq \psi(\mathbb{B})$ and $\eta \in \overline{S}$. According to [7, lemma 2.21] $\mathcal{O}_{X,\eta}$ is a field because $\mathbb{B}^2$ is a neighbourhood of η in $\mathbb{A}_k^{2,an}$. Furthermore $\overset{\circ}{S} = \emptyset$ because $S \subsetneq Z := \{x \in \mathbb{B}_{(r,1)} \mid T_2(x) = f(T_1(x))\}$, which is a Zariski closed subset of dimension 1 of $\mathbb{B}_{(r,1)}$, which itself is of pure dimension 2, so Z is nowhere dense in $\mathbb{B}_{(r,1)}$ [1, 2.3.7]. Hence according to corollary 2.3, S is not overconvergent constructible. However, if we consider the covering of X

given by $X_1 = \{x \in X \mid |T_1(x)| \leq r\}$, $X_2 = \{x \in X \mid |T_1(x)| \geq r\}$, then $S \cap X_1$ is indeed globally semianalytic in X_1 and $S \cap X_2 = \emptyset$, so S is rigid semianalytic.
Now since the class of overconvergent subanalytic subsets contains the class of globally semianalytic subsets, if the class of overconvergent subanalytic subsets was G-local, it should contain the class of rigid semianalytic subset, but we have shown that this is false. Hence the class of overconvergent subanalytic subsets is not G-local. ☐

Remark 2.5. Actually, this example gives directly a counterexample to [19, lemma 4.1] which in our feeling is the source of mistakes in [19].

As a corollary of this we obtain:

Proposition 2.6. *Let $0 < s < r < 1$ with $s, r \in \sqrt{|k^\times|}$, $f \in k\{r^{-1}u\}$ whose radius of convergence is exactly r such that $\|f\| < 1$, and let's set $\mathbb{B}^2 = \mathcal{M}(k\{T_1, T_2\})$. Define :*

$$S = \{x \in \mathbb{B}^2 \mid |T_1(x)| \leq s \text{ and } T_2(x) = f(T_1(x))\}$$

Then S is a locally semianalytic subset of $\mathbb{B}^2$ which is not a globally semianalytic subset of $\mathbb{B}^2$.

Proof. If S was a globally semianalytic subset of $\mathbb{B}^2$, we could find $T \subseteq S$ which contains infinitely many points of S such that T is a basic semianalytic subset, and even, a finite intersection of sets of the form $\{x \in \mathbb{B}^2 \mid |g_1(x)| < |g_2(x)|\}$, $\{x \in \mathbb{B}^2 \mid |g_1(x)| \leq |g_2(x)| \neq 0\}$ and $\{x \in \mathbb{B}^2 \mid h(x) = 0\}$. Since an intersection of the two first kind of sets is a strictly analytic domain, and $T \subseteq S$, and $\overset{\circ}{S} = \emptyset$, in this intersection, there must be a non-trivial set of the form $\{x \in \mathbb{B}^2 \mid h(x) = 0\}$. Now, let's consider in $\mathbb{B}_{(r,1)} = \mathcal{M}(k\{r^{-1}T_1, T_2\})$ the Zariski-closed subset $Z = V(T_2 - f(T_1), h)$. By assumption, it is infinite. Moreover, since $\|f\| < 1$, $T_2 - f(T_1)$ is irreducible (see the lemma above) in $\mathcal{M}(k\{r^{-1}T_1, T_2\})$, so for reasons of dimension, in $\mathcal{M}(k\{r^{-1}T_1, T_2\})$, $V(T_2 - f(T_1)) \subseteq V(h)$. But now if we introduce (as in the preceding proof)

$$\begin{array}{ccc} \mathbb{B}_r & \xrightarrow{\psi} & \mathbb{B}^2 \\ u & \mapsto & (u, f(u)) \end{array}$$

and $\eta = \psi(g)$ where g is the Gauss point of $\mathbb{B}_r$, then $\eta \in V(h)$ (where we see now $V(h)$ as a Zariski closed subset of $\mathbb{B}^2$), $\mathcal{O}_{\mathbb{B}^2,\eta}$ is a field, but $\overset{\circ}{V(h)} = \emptyset$, and since $V(h)$ is a globally semianalytic (so overconvergent constructible) subset of $\mathbb{B}_{(r,1)}$, this contradicts corollary 2.3.
Let us now show that S is locally semianalytic subset of $\mathbb{B}^2$. Indeed, take $0 < s < t < r$, and consider $X_1 = \{x \in \mathbb{B}^2 \mid |T_1(x)| \leq r\}$ and $X_2 = \{x \in \mathbb{B}^2 \mid |T_1(x)| \geq t\}$. They define a wide covering of $\mathbb{B}^2$ and $X_1 \cap S$ (resp. $X_2 \cap S$) is globally semianalytic in X_1 (resp. X_2), so S is well locally semianalytic in $\mathbb{B}^2$. ☐

We've implicitly used:

Lemma 2.7. *If $f \in k\{r^{-1}x\}$ and $\|f\| \leq 1$, then $F(x, y) := y - f(x)$ is irreducible in $k\{r^{-1}x, y\}$.*

Proof. As we have already seen, $V(f)$ is isomorphic to $\mathbb{B}_r$, so is irreducible. ☐

2.3. **The other inequalities.**

We'll now explain the other inequalities appearing in figure 1.
The following proposition will be implicitly used in the rest of this section. In addition, it illustrates that the mixture of overconvergence and rigid semianalytic subsets (which is a G-local property), is somehow too strong, in the sense that in proposition 2.8 above, the overconvergence condition seems to have disappeared.

Proposition 2.8. *Let $S \subseteq X$. The following properties are equivalent:*

1. *S is strongly subanalytic.*
2. *There exist $n \in \mathbb{N}$ and $T \subseteq X \times \mathbb{B}^n$ a rigid semianalytic subset such that*

$$S = \pi(T \cap (X \times (\overset{\circ}{\mathbb{B}})^n))$$

where $\pi : X \times \mathbb{B}^n \to X$ is the natural projection.

Proof. Let's show that (1) $\Rightarrow$ (2). Let S be a strongly subanalytic subset of X, so there exists $r > 1$, $T \subseteq X \times \mathbb{B}^n_r$ a rigid semianalytic subset such that $S = \pi(T \cap (X \times \mathbb{B}^n))$. Decreasing r if necessary, we can assume that $|r| \in \sqrt{|k^*|}$. In fact, using similar arguments as the one given in remark 1.15, we can even assume that $r \in |k|$. Then if we consider the homothety, which is an isomorphism: $h : X \times \mathbb{B}^n_r \to X \times \mathbb{B}^n$, which can be defined as multiplication of each coordinate of $\mathbb{B}^n_r$ by $\frac{1}{\lambda}$, this gives the following commutative diagram:

$$\begin{tikzcd}
X \times \mathbb{B}^n_r \arrow[r, "h"] \arrow[dr, "\pi"'] & X \times \mathbb{B}^n \arrow[d, "\pi'"] \\
 & X
\end{tikzcd}$$

and $S = \pi(T \cap (X \times \mathbb{B}^n) = \pi'\left(h(T) \cap (X \times \mathbb{B}^n_{\frac{1}{r}})\right)$. Now $T' := h(T) \cap (X \times \mathbb{B}^n_{\frac{1}{r}})$ is a rigid semianalytic subset of $X \times \mathbb{B}^n$) such that $T' \subseteq X \times (\overset{\circ}{\mathbb{B}})^n$ and $S = \pi'(T') = \pi'(T' \cap (X \times (\overset{\circ}{\mathbb{B}})^n))$.

Conversely, let $T \subseteq X \times (\overset{\circ}{\mathbb{B}})^n$ be a rigid semianalytic subset of $X \times \mathbb{B}^n$ and $S = \pi(T)$. For any $r > 1$, we can define $X_0 = X \times \mathbb{B}$, and for $i = 1 \dots n$, let $X_i = \{(x, t_1, \dots, t_n) \in X \times \mathbb{B}^n_r \mid |t_i| \geq 1\}$. So $\{X_i\}_{i \in \{0 \dots n\}}$ is an admissible covering of $X \times \mathbb{B}^n_r$. By assumption, $T \cap X_0$ is rigid semianalytic, and $T \cap X_i = \emptyset$ for $i = 1 \dots n$. So T is rigid semianalytic in $X \times \mathbb{B}^n_r$, and if we note $\pi : X \times \mathbb{B}^n_r \to X$, $S = \pi(T)$, so S is strongly subanalytic. ☐

Proposition 2.9. *There exist strongly subanalytic subsets which are not rigid overconvergent constructible.*

Proof. Let $r > 1$, $X = \mathcal{M}(k\{x, y, z\}) = \mathbb{B}^3$, $Y = \mathcal{M}(k\{x, y, z, t\})$ and $\pi : Y = \mathcal{M}(k\{x, y, z, t\}) \to X = \mathcal{M}(k\{x, y, z\})$, the natural projection. We now choose $f \in k\{t\}$ whose radius of convergence is exactly 1, and such that $\|f\| \leq 1$, and $T = \{(x, y, z, t) \in Y \mid |t| < 1, x = yt, z = yf(t)\}$. It is a rigid semianalytic subset of Y, and $S = \pi(T)$ is a strongly subanalytic subset of X according to the previous proposition. Since the family of closed balls with center the origin is a fundamental system of neighbourhoods of the origin, if S was rigid overconvergent constructible, for some $1 \geq |\mu| = \varepsilon > 0$ small enough, $S' := S \cap \mathbb{B}^3_\varepsilon$ would be overconvergent constructible in $\mathbb{B}^3_\varepsilon$. We then fix a $y_0 \in k^*$ such that $0 < |y_0| < \varepsilon$, i.e. $\frac{|y_0|}{|\mu|} < 1$ and define $X' := \{(x, y, z) \in \mathbb{B}^3_\varepsilon \mid y = y_0\}$. Now X' is isomorphic to the bidisc $\mathbb{B}^2_\varepsilon = \{(x, y) \mid |x| \leq \varepsilon \text{ and } |y| \leq \varepsilon\}$, and $S'' := S \cap X'$ should be overconvergent constructible in X' (somehow we use here lemma 1.41 (2)). If we make a dilatation of X' by $\frac{1}{\mu}$ it becomes the bidisc of radius 1: the new coordinates are x', z' defined by $x = \mu x'$ and $z = \mu z'$. Now, in these new coordinates:

$$S'' = \{(x', z') \in \mathbb{B}^2 \mid |x'| < \frac{|y_0|}{|\mu|} \text{ and } z' = \frac{y_0}{\mu} f(\frac{x'\mu}{y_0})\}$$

should be overconvergent constructible in $\mathbb{B}^2$. If we put $r := \frac{|y_0|}{|\mu|} < 1$ and $g(x') = \frac{y_0}{\mu} f(\frac{x'\mu}{y_0})$, then the radius of convergence of g is precisely r, $\|g\| < 1$ so $S'' = \{(x', z') \in \mathbb{B}^2 \mid |x'| < r \text{ and } z' = g(x')\}$, S'' should be overconvergent constructible in $\mathbb{B}^2$, but we proved the converse in proposition 2.4. ☐

Proposition 2.10. *There are overconvergent constructible subsets which are not rigid semianalytic.*

Proof. Let $1 < r = |\lambda|$, and $f \in k\{r^{-1}X\}$ whose radius of convergence is exactly r, and such that $\|f\| < 1$. We set $X = \mathbb{B}^3 = \mathcal{M}(k\{x, y, z\})$, $Y = \mathcal{M}(k\{x, y, z, r^{-1}t\})$, and

$$T = \{(x, y, z, t) \in Y \mid x = yt,\ z = yf(t),\ |t| \leq 1\}$$

and $S = \pi(T)$, where $\pi : \mathcal{M}(k\{x, y, z, r^{-1}t\}) \to \mathcal{M}(k\{x, y, z\})$ is the natural projection. S is then overconvergent subanalytic, so overconvergent constructible. If S was rigid semianalytic, there would exist $\mu \in k$, with $0 < \varepsilon := |\mu| < 1$ such that $S' = S \cap \mathbb{B}^3$ is globally semianalytic in $\mathbb{B}^3_\varepsilon$ (we *again* use that if V is an affinoid domain of $\mathbb{B}^3$ that contains the origin, then there exists $\varepsilon > 0$ such that $\mathbb{B}^3_\varepsilon \subseteq V$). Let us introduce $y_0 \in k^*$ such that $0 < |y_0| < \frac{\varepsilon}{r}$. In particular $\frac{|y_0|}{\varepsilon} = \left|\frac{y_0}{\mu}\right| < \frac{1}{r}$.

Then $X' = \{(x,y,z) \in \mathbb{B}^3_\varepsilon \mid y = y_0\}$ is a Zariski-closed subset of $\mathbb{B}^3_\varepsilon$, isomorphic to a bidisc $\mathbb{B}^2$. Now, $S'' := S \cap X'$ is defined by

$$S'' = \{(x,z) \in \mathbb{B}^2_\varepsilon \mid \left|\frac{x}{y_0}\right| \leq 1 \text{ and } z = y_0 f(\frac{x}{y_0})\}$$

As we said, X' is isomorphic to $\mathbb{B}^2$ with coordinates (x',z') where $x = \mu x'$ and $z = \mu z'$. In these new coordinates, $S'' = \{(x',z') \in \mathbb{B}^2 \mid \left|\frac{x'\mu}{y_0}\right| \leq 1 \text{ and } z'\mu = y_0 f(\frac{x'\mu}{y_0})\}$. If we define $g(x') = \frac{y_0}{\mu} f(\frac{x'\mu}{y_0})$ and $s = \frac{|y_0|}{\varepsilon} = \left|\frac{y_0}{\mu}\right| < \frac{1}{r}$, then g has a radius of convergence which is exactly ρ where $s < \rho = \left|\frac{y_0}{\mu}\right| r < 1$, and $\|g\| < \|f\| < 1$, so $S'' = \{(x',z') \in \mathbb{B}^2 \mid |x'| \leq s \text{ and } z' = g(x')\}$, should be globally semianalytic, but is not (see proposition 2.6). ☐

From this one can deduce:

Corollary 2.11. *Let X be a k-analytic space which contains a closed ball of dimension ≥ 3. Then there are overconvergent subanalytic subsets of X which are not rigid semianalytic. In particular, the class of overconvergent subanalytic subsets of X properly contains the class of locally semianalytic subsets of X.*

In conclusion, in figure 1, we have shown inequalities $1, 4, 5$ and 8. Now $2, 3, 6, 7$ are set-theoretical consequences of 4, 5 and of the inclusions from the left to the right.

2.4. **Berkovich points versus rigid points.** Let $X = \mathcal{M}(\mathcal{A})$. Talking about globally semianalytic or overconvergent constructible subsets S of X, one can wonder if things change if we restrict to $S_{rig} = S \cap X_{rig}$. Actually the following two propositions show that there is no difference if one works with Berkovich spaces or rigid spaces. To be precise, let $\mathcal{B}$ be the free boolean algebra whose set of variables consists in the set of *formal inequalities* $\{|f| \leq |g|\}$, $\{|f| < |g|\}$ and $\{f = 0\}$, for $f, g \in \mathcal{A}$. If we note GSA_{rig} the class of globally semianalytic subsets of X_{rig} and GSA_{Ber} the class of globally semianalytic subset of the Berkovich space X, we have a commutative diagram, where for instance $\alpha(\{|f| \leq |g|\}) = \{x \in X \mid |f(x)| \leq |g(x)|\}$ and $\beta(\{|f| \leq |g|\}) = \{x \in X_{rig} \mid |f(x)| \leq |g(x)|\}$:

$$\begin{tikzcd}
\mathcal{B} \arrow[r, "\alpha"] \arrow[dr, "\beta"'] & GSA_{Ber} \arrow[d, "\iota"] \\
 & GSA_{rig}
\end{tikzcd}$$

Proposition 2.12. *In the above figure, ι is bijective.*

Proof. First, ι is surjective because by definition β is surjective. Now if $\iota(S_1) = \iota(S_2)$, we must show that $S_1 = S_2$. Considering $S_1 \setminus S_2$ and $S_2 \setminus S_1$, we can restrict to show that if $S \in GSA_{Ber}$ and $\iota(S) = \emptyset$, then $S = \emptyset$. According to what has been previously done, we can assume that $S \in GSA_{Ber}$ is a finite intersection of subsets of the form $\{x \in X \mid |f(x)| \leq |g(x)| \neq 0\}$, $\{x \in X \mid |f(x)| < |g(x)|\}$ and $\{x \in X \mid h(x) = 0\}$, and that $\iota(S) = S \cap X_{rig} = \emptyset$. Passing to $Y = \mathcal{M}(\mathcal{A}/\mathcal{I})$ where $\mathcal{I}$ is the ideal generated by the functions h appearing in the third case ($h(x) = 0$), we can assume that S is a finite intersection of subsets of the form: $\{x \in X \mid |f(x)| \leq |g(x)| \neq 0\}$ and $\{x \in X \mid |f(x)| < |g(x)|\}$. But then it forms a non empty strictly analytic domain of X so $S \cap X_{rig} \neq \emptyset$. ☐

If we denote by CD the family of finite subsets of constructible data of X, by OC the family of overconvergent constructible subsets of X, and OC_{rig} the family of subsets of X_{rig} which are the intersection of an element of OC with X_{rig}, (as we proved, in [19] these are the globally strongly subanalytic, or globally strongly D-semianalytic subsets of X_{rig}), then we have the following commutative diagram:

$$\begin{tikzcd}
CD \arrow[r, "\alpha"] \arrow[dr, "\beta"'] & OC \arrow[d, "\iota"] \\
 & OC_{rig}
\end{tikzcd}$$

To be precise, if $\mathcal{D} \in CD$ is the set of the constructible data $(X_i, T_i) \overset{\varphi_i}{\dashrightarrow} X$, then

$$\alpha(\mathcal{D}) = \bigcup_{i=1}^{n} \varphi_i(T_i).$$

Proposition 2.13. *In the above digram, ι is a bijection.*

Proof. Since we showed that OC (and OC_{rig}) is stable under complementary, we can restrict to show that if $S \in OC$ is such that $\iota(S) = S \cap X_{rig} = \emptyset$, then $S = \emptyset$. To show this we can even assume that $S = \varphi(T)$, where $(Y, T) \overset{\varphi}{\dashrightarrow} X$ is a constructible datum. But, if T is a non empty globally semianalytic subset of Y, according to proposition 2.12, $T_{rig} \neq \emptyset$, so since φ preserves the rigid points, $\varphi(T)_{rig} = S_{rig}$ is non empty.

□

3. Overconvergent subanalytic subsets when $\dim(X) = 2$

In this section, k will be non-archimedean algebraically closed field. In this context, if X is a k-analytic space, it is equivalent to say that X is regular, or that X is quasi-smooth [7, section 3]. We'll rather use the second terminology.

Let X, Y be two k-analytic spaces, so that we can consider the cartesian diagram

$$\begin{tikzcd} & X \times Y \arrow[dl, "\pi_1"'] \arrow[dr, "\pi_2"] & \\ X & & Y \end{tikzcd}$$

Let $z \in X \times Y$, and let's denote by $z_1 = \pi_1(z)$, $z_2 = \pi_2(z)$. We'll assume that $z_2 \in Y(k) = Y_{rig}$.

Lemma 3.1. *Let V be an affinoid domain of $X \times Y$ such that $z \in V$. There exists an affinoid domain U of X (which contains z_1) such that if W is an affinoid neighbourhood of z_2 small enough, $V \cap (X \times W) = U \times W$.*

Proof. [20, 2.2] □

Corollary 3.2. *Let $\mathcal{V}$ be a neighbourhood of z. There exists U (resp. W) an affinoid neighbourhood of z_1 (resp. z_2) such that $\mathcal{V} \supseteq U \times W$*

Proof. We can assume that $\mathcal{V} = V$ is an affinoid neighbourhood. In the previous lemma, $V \cap (X \times W)$ is still a neighbourhood of z, since W is an affinoid neighbourhood of z_2 (because z_2 is a rigid point). If we note $s_{z_2} : X \to X \times Y$ the section of π_1 defined by $s_{z_2}(t) = (t, z_2)$, then $s_{z_2}^{-1}((V \cap (X \times W)) = s_{z_2}^{-1}(U \times W) = U$ is an affinoid neighbourhood of x (since $s_{z_2}(x) = z$). □

Remark 3.3. Without the assumption that $z_2 \in Y(k)$ the previous corollary would be false. Indeed assume that k is algebraically closed. Take for instance $X = \mathcal{M}(k\{x\})$ and $Y = \mathcal{M}(k\{y\})$, and let $\varphi : \mathcal{M}(k\{t\}) \to X \times Y$ be defined by $\varphi(t) = (t, -t)$. Let η be the Gauss point of $\mathcal{M}(k\{t\})$ and $z := \varphi(\eta)$. Let $V = \{p \in \mathcal{M}(k\{x, y\}) \mid |(x+y)(p)| \leq \frac{1}{2}\}$. It is a neighbourhood of z. However, $\pi_1(z)$ (resp $\pi_2(z)$) is the Gauss point $z_1 = \eta_X$ of $\mathcal{M}(k\{x\})$ (resp. $z_2 = \eta_Y$ the Gauss point of $\mathcal{M}(k\{y\})$). It is then easy to see, according to the description of affinoid domain of the unit disc as Swizz cheese, that there doesn't exist an affinoid neighbourhood U (resp. W) of η_X (resp. η_Y) such that $V \supseteq U \times W$. For instance for the reason that in U there would necessarily exist a rigid point $x_0 \in k^0$ such that $\overline{x_0} = \overline{0}$ and in W a rigid point y_0 such that $\overline{y_0} = \overline{1}$ but $(x_0, y_0) \notin V$ (where $\overline{x}$ corresponds to the reduction of x in $\tilde{k}$).

Lemma 3.4. *Let $x \in X = \mathcal{M}(\mathcal{A})$, and $f = \sum_{n \in \mathbb{N}} a_n T^n \in \mathcal{A}\{r^{-1}T\}$ such that $f_x \neq 0$. Then there exists an affinoid domain $V = \mathcal{M}(\mathcal{B})$ which contains x, $P \in \mathcal{B}[T]$, and $u \in \mathcal{B}\{r^{-1}T\}$ a multiplicative unit such that $f_{|V \times \mathbb{B}_r} = (uP)_{|V \times \mathbb{B}_r}$.*

Proof. Since $f_x = \sum_{n \in \mathbb{N}} a_n(x) T^n \neq 0$, this series is distinguished of some order $s \geq 0$, which means that $|a_s(x)| r^s = \|f_x\|$ and that s is the greatest rank for this property.

We now use lemme 1.27 in our specific situation where the polyradius $\underline{r}$ is in fact a real number r. Hence we can introduce a finite subset $J \subseteq \mathbb{N}$ and some series $\phi_n \in \mathcal{A}\{r^{-1}T\}$ for $n \in J$ satisfying

$\|\phi_n\| < r^n$ such that $f = \sum\limits_{n\in J} a_n(X^n + \phi_n)$. One checks that $s \in J$.
We then define V as the rational domain:

$$V = \{z \in X \mid |a_s(z)| = |a_s(x)| \text{ and } |a_i(z)|r^i \leq |a_s(x)|r^s \ i \in J \setminus \{s\}\}.$$

It is true that $x \in V$. Moreover, on $V = \mathcal{M}(\mathcal{B})$, one checks that a_s is a multiplicative unit, and that on $\mathcal{B}\{r^{-1}T\}$, f is distinguished of order s. One can then apply Weierstrass preparation (corollary 1.26) to conclude. □

Remark 3.5. The previous result (lemma 3.4) is false if we remove the assumption $f_x \neq 0$: Indeed, if $0 < r < 1$, $f \in k\{r^{-1}x\}$ whose radius of convergence is exactly r and such that $\|f\| < 1$. Let then $\mathcal{A} = k\{y,t\}$, $X = \mathcal{M}(\mathcal{A})$ the unit bidisc, p the rigid point of X corresponding to the origin, and $F(x,t,y) = y - tf(x) \in k\{y,t\}\{r^{-1}x\} = \mathcal{A}\{r^{-1}X\}$. Then there does not exist an affinoid domain $V = \mathcal{M}(\mathcal{B})$ containing p such on $V \times \mathbb{B}_r$, $F_{|V\times\mathbb{B}_r} = (e.P)_{|V\times\mathbb{B}_r}$ where e is a multiplicative unit and $P \in \mathcal{B}[t]$.

Proof. Indeed otherwise, we could assume that V is the closed bidisc of radius say $|\lambda| \in |k^\times|$ and then fixing $t = \lambda$, we could conclude that if we consider $G = y - \lambda f(x) \in k\{r^{-1}x, y\}$, there exist $P \in k\{y\}[x]$ and e a multiplicative unit of $k\{x,y\}$ such that $G = eP$. But then this would imply that if we set

$$S := \{(x,y) \in \mathbb{B}^2 \mid |x| \leq r \text{ and } y = f(x)\}$$

then $S = \{(x,y) \in \mathbb{B}^2 \mid |x| \leq r \text{ and } P(x,y) = 0\}$ so that S would be globally semianalytic in $\mathbb{B}^2$, but in section 2, we exploited many times that this is not the case. □

From now on, X will be a regular Berkovich space of dimension 2.
We know make two simple remarks that we'll used in the proof of theorem 3.13.

Lemma 3.6. *Let $\mathcal{A}$ be an affinoid algebra, $X = \mathcal{M}(\mathcal{A})$, $0 < r < s$ some real numbers and $h \in \mathcal{A}$.*

1. *Consider the Weierstrass domain of X: $V = \{x \in X \mid |h(x)| \leq s\}$ and let S be a locally semi-analytic subset of V such that $S \subseteq \{x \in X \mid |h(x)| \leq r\}$. Then S, seen as a subset of X, is a also a locally semianalytic subset of X.*
2. *Consider the Laurent domain of X: $V = \{x \in X \mid |h(x)| \geq r\}$ and let S be a locally semi-analytic subset of V such that $S \subseteq \{x \in X \mid |h(x)| \geq s\}$. Then S, seen as a subset of X, is a also a locally semianalytic subset of X.*

Proof. Choose a real number t such that $r < t < s$.

1. Set $W = \{x \in X \mid |h(x)| \geq t\}$. Then $\{V, W\}$ is a wide covering of X, and $S \cap V$ is by hypothesis locally semianalytic in V, and by assumption, $S \cap W = \emptyset$ so is also locally semianalytic in W, hence S is locally semianalytic in X.
2. Likewise, set $W = \{x \in X \mid |h(x)| \leq t\}$. Then $\{V, W\}$ is a wide covering of X, $S \cap V$ is locally semianalytic in V and $S \cap W = \emptyset$, so S is locally semianalytic in X.

□

This lemma will be used jointly with the following remark:

Remark 3.7. Let's consider the following situation: $X = \mathcal{M}(\mathcal{A})$, $f, g \in \mathcal{A}$, $0 < s < r$ and $(Z,S) \xrightarrow{\varphi}$ the elementary constructible datum given by $Z = \mathcal{M}(\mathcal{B})$ where $\mathcal{B} = \mathcal{A}\{r^{-1}t\}/(f - tg)$ and $S = \{z \in Z \mid g(z) \neq 0 \text{ and } |f(z)| \leq s|g(z)|\}$. Moreover, let $(Y,U) \overset{\psi}{\dashrightarrow} (Z,S)$ be a constructible data.

1. If $g|f$, there exists $h \in \mathcal{A}$ such that $f = gh$. Let then $\mathcal{C} = \mathcal{A}\{r^{-1}t\}/(h - t)$, $V = \mathcal{M}(\mathcal{C})$. Note that V is the Weierstrass domain of X defined by $V = \{x \in X \mid |h(x)| \leq r\}$. Let's denote by β the map of the immersion of the affinoid domain V inside X, and let $T = \{x \in V \mid |h(x)| \leq s\}$. Since $f - tg = g(h - t)$, $(h - t)|(f - tg)$, and there is a closed immersion $V \xrightarrow{\alpha} Z$. Moreover, $\alpha(T) = S$.
Indeed $\alpha(T) \subseteq S$, follows from their respective definitions. Conversely, if $z \in S$, $(f - tg)(z) = 0 = g(z)(h - t)(z)$ but since $g(z) \neq 0$, $(h - t)(z) \neq 0$ which implies that $z \in V$,

and by the definition of S, it follows that $z \in \alpha(T)$.
Let's then consider the following cartesian diagram of k-germs:

$$\begin{tikzcd}
(Y,U) \arrow[r, dotted, "\psi"] & (Z,S) \arrow[r, "\varphi"] & X \\
(Y',U') \arrow[u, "\alpha'"] \arrow[r, dotted, "\psi'"] & (V,T) \arrow[u, "\alpha"] \arrow[ur, "\beta"] &
\end{tikzcd}$$

Here, $(Y',U') \overset{\psi'}{\dashrightarrow} (V,T)$ is still a constructible datum. Since $\alpha(T) = S$, it follows that $\alpha(\psi'(U'))) = \psi(U)$, so

$$\varphi(\psi(U)) = \varphi(\alpha(\psi'(U'))) = \beta(\psi'(U')). \tag{11}$$

Roughly speaking, we were starting with the constructible datum $(Y,U) \overset{\psi}{\dashrightarrow} (Z,S) \xrightarrow{\varphi} X$ such that the elementary constructible datum of φ was defined with f, g such that $g|f$. And we have been able to replace it by the following constructible datum $(Y',U') \overset{\psi'}{\dashrightarrow} (V,T) \xrightarrow{\beta} X$ where V is a Weierstrass domain. Note moreover that T and so also $\psi'(U')$ satisfy the hypothesis of lemma 3.6 (1).

(2) If $f|g$, there exists $h \in \mathcal{A}$ such that $g = fh$. Let then $\mathcal{C} = \mathcal{A}\{r^{-1}t\}/(1-th)$, $V = \mathcal{M}(\mathcal{C})$. Note that V is the Laurent domain of Xdefined by $V = \{x \in X \mid |h(x)| \geq \frac{1}{r}\}$. Let's denote by β the map of the immersion of the Laurent domain V inside X, and let $T = \{x \in V \mid |h(x)| \geq \frac{1}{s}\}$. Since $(1-th)|(f-tg)$, there is a closed immersion $V \xrightarrow{\alpha} Z$. Moreover, $\alpha(T) = S$.
We then consider the following cartesian diagram of k-germs:

$$\begin{tikzcd}
(Y,U) \arrow[r, dotted, "\psi"] & (Z,S) \arrow[r, "\varphi"] & X \\
(Y',U') \arrow[u, "\alpha'"] \arrow[r, dotted, "\psi'"] & (V,T) \arrow[u, "\alpha"] \arrow[ur, "\beta"] &
\end{tikzcd}$$

Here, $(Y',U') \overset{\psi'}{\dashrightarrow} (V,T)$ is still a constructible datum. Since $\alpha(T) = S$, it follows that $\alpha(\psi'(U'))) = \psi(U)$, so

$$\varphi(\psi(U)) = \varphi(\alpha(\psi'(U'))) = \beta(\psi'(U')). \tag{12}$$

In that case, we were starting with the constructible datum $(Y,U) \overset{\psi}{\dashrightarrow} (Z,S) \xrightarrow{\varphi} X$ such that $f|g$, and we have been able to replace it by the following constructible datum $(Y',U') \overset{\psi'}{\dashrightarrow} (V,T) \xrightarrow{\beta} X$ where V is a Laurent domain. Note moreover that T and so also $\psi'(U')$ satisfies the hypothesis of lemma 3.6 (2).

Remark 3.8. We are going to use some blowing up of k-analytic spaces in the following context: X will be a quasi-smooth k-analytic space of dimension 2, and we'll blow up a rigid point p of X. In particular, the resulting blowing up $\tilde{X}$ will be still quasi-smooth. To give a precise description of the situation, since k is algebraically closed, we can assume that $X = \mathbb{B}^2$ and p is the origin. The blowing up can then be described with two charts as follows. We consider

$$\begin{array}{rcc|rcc}
X_1 = \mathcal{M}(k\{x,t_1\}) & \xrightarrow{\pi_1} & \mathbb{B}^2 = \mathcal{M}(k\{x,y\}) & X_2 = \mathcal{M}(k\{y,t_2\}) & \xrightarrow{\pi_2} & \mathbb{B}^2 = \mathcal{M}(k\{x,y\}) \\
(x,t_1) & \mapsto & (x,t_1x) & (y,t_2) & \mapsto & (t_2y,y)
\end{array}$$

Then $\tilde{\mathbb{B}^2}$ is obtained by gluing X_1 and X_2 along the domains $U_1 = \{z \in X_1 \mid t_1(z) \neq 0\}$ and $U_2 = \{z \in X_2 \mid t_2(z) \neq 0\}$ via the isomorphism

$$\begin{array}{ccc}
U_1 & \to & U_2 \\
(x,t_1) & \mapsto & (xt_1, t_1^{-1}).
\end{array}$$

Proposition 3.9. *Let $X = \mathcal{M}(\mathcal{A})$ be a quasi-smooth k-analytic space of dimension 2 and $f, g \in \mathcal{A}$. Then there exists a succession of blowing up of rigid points $\pi : \tilde{X} \to X$ such that for all $x \in \tilde{X}$, $f_x|g_x$ or $g_x|f_x$. Note that $\tilde{X}$ is quasi-smooth.*

Proof. We may assume that X is irreducible. If $f = 0$ or $g = 0$, there is nothing to prove, so we may assume that $f \neq 0$ and $g \neq 0$. Likewise, if $f = g$, there is nothing to do, so we may also assume that $f - g \neq 0$
Let $h = fg(f-g)$. Hence, $h \neq 0$. We can find a succession of blowing up of rigid points $\pi : \tilde{X} \to X$ such that $\pi^*(h)$ is a normal crossing divisor. Indeed, the classical proof (see [14, 1.8]) that works in the algebraic case, or the complex analytic case, can be translated *verbatim* in our context, and since we are dealing with a compact space, the local procedure needs only to be applied to a finite number of points.
Let then $x \in \tilde{X}$. If it is not a rigid point, $\mathcal{O}_{\tilde{X},x}$ is a field or a discrete valuation ring and the result is clear.
Otherwise $h = fg(f-g)$ is a normal crossing divisor, and it can be written in $\mathcal{O}_{\tilde{X},x}$ as $u\xi_1^n\xi_2^m$ where ξ_1, ξ_2 is a system of local parameters around x and u is a unit in $\mathcal{O}_{\tilde{X},x}$. Dividing by the common divisor of f and g in $\mathcal{O}_{\tilde{X},x}$, we can assume for instance that $f = v\xi_1^p$ and $g = w\xi_2^q$ and $f - g = z\xi_1^a\xi_2^b$ where v, w and z are units. If $p > 0$ then modulo ξ_1 we obtain $f = 0$, so $f - g = w\xi_2^q$ modulo ξ_1. This implies that $a = 0$ and $b = q$. So $f = (f-g) + g$ is divisible by ξ_2^q, in fact $q = 0$, and $g_x | f_x$.

□

Lemma 3.10 (Local algebraization of a function in a family of rings)**.** *Let* $Y \subseteq \mathcal{M}(k\{T\})$ *be the Laurent domain* $Y = \{y \in \mathcal{M}(k\{T\}) \mid |(T-a_0)(y)| \leq r_0$ *and* $|(T-a_i)(y)| \geq r_i,\ i = 1 \dots n\}$*, and let* $X = \mathcal{M}(\mathcal{A})$ *be an affinoid space. Let* $f \in \mathcal{O}(X \times Y)$*,* $z \in X \times Y$*, such that* $\pi_1(z) = x \in X(k)$ *and set* $y := \pi_2(z)$*. Assume that* $f_x \in \mathcal{H}(x) \otimes \mathcal{O}(Y) \simeq \mathcal{O}(Y)$ *is non-zero*[4]*. Then there exists* $V = \mathcal{M}(\mathcal{B})$ *an affinoid neighbourhood of* x*,* $Y' = \{y \in \mathcal{M}(k\{T\}) \mid |(T-b_0)(y)| \leq s_0$ *and* $|(T-b_i)(y)| \geq s_i,\ i = 1 \dots m\}$ *an affinoid neighbourhood of* y *such that* $f_{|V\times Y'} = (u.P)_{|V\times Y'}$ *where* u *is a multiplicative unit of* $V \times Y'$ *and* $P \in \mathcal{B}[T, (T-b_1)^{-1}, \dots, (T-b_m)^{-1}]$.

Remark 3.11. Let's mention that in the proof we distinguish two very different cases:

1. If y is a rigid point then Y' can in fact be chosen to be a closed ball, i.e. $m = 0$.
2. Otherwise, if y is not a rigid point, then in fact $s_0 = r_0$, that is to say, we don't have to decrease the radius of the ambient closed ball, but in counterpart, we possibly have to remove some open balls.

Proof. If y is a rigid point, we can indeed find a closed disc Y' which contains y and the result follows from lemma 3.4.
If y is not a rigid point, $f_x \in \mathcal{H}(x) \otimes \mathcal{O}(Y) \simeq \mathcal{O}(Y)$. Then (cf [9, 2.2.9]) there exist $\alpha_1, \dots, \alpha_N \in k$, $d_1, \dots, d_N \in \mathbb{N}$, such that $f_x = \prod_{i=1}^N (T-\alpha_i)^{d_i} g$ where g is an invertible function of $\mathcal{O}(Y)$. We then set $m = n + N$, $b_0 = a_0$, $b_i = a_i$, $s_i = r_i$ for $i = 1 \dots n$, and $b_{n+j} = \alpha_j$ for $j = 1 \dots N$ and s_{n+j} small enough so that $\{|T - \alpha_j| \geq s_{n+j}\}$ is a neighbourhood of y (this is possible because y is not a rigid point). Define

$$Y' = \{y \in \mathcal{M}(k\{T\}) \mid |(T-b_0)(y)| \leq s_0 \text{ and } |(T-b_i)(y)| \geq s_i, i = 1 \dots m\}$$

We then set $G = f\prod_{i=1}^N (T-\alpha_i)^{-d_i} \in \mathcal{O}(X \times Y')$. Then $G_x = g$ doesn't vanish on Y'_x, so there exists an affinoid neighbourhood $V = \mathcal{M}(\mathcal{B})$ of x such that G is invertible on $V \times Y'$ because the locus of points x where G_x is invertible is open. Now using the explicit description of $\mathcal{O}(V \times Y')$,

$$G = \sum_{\nu=(\nu_0,\dots,\nu_n)\in\mathbb{N}^{m+1}} b_\nu (T-b_0)^{\nu_0}(T-b_1)^{-\nu_1}\dots(T-b_m)^{-\nu_m} \tag{13}$$

Now for $M \geq 0$ set

$$G_M = \sum_{|\nu|\leq M} b_\nu (T-b_0)^{\nu_0}(T-b_1)^{-\nu_1}\dots(T-b_m)^{-\nu_m} \tag{14}$$

By definition, $G_M \in \mathcal{B}[T, (T-b_1)^{-1}, \dots, (T-b_m)^{-1}]$, $G_M \xrightarrow[M\to\infty]{} G$, so G_M is invertible for M big enough, and for such a M, $G = (G - G_M) + G_M = G_M(1 + G_M^{-1}(G - G_M))$, and $\|G_M^{-1}\| = \|G^{-1}\|$. Now:

$$\|G_M^{-1}(G - G_M)\| \xrightarrow[M\to\infty]{} 0$$

---

[4] Here $\mathcal{H}(x) \simeq k$ because $x \in X(k)$.

so $u_M = 1 + G_M^{-1}(G - G_M)$ is a multiplicative unit for M big enough, and

$$f = G_M u_M \prod_{i=1}^{N} (T - \alpha_i)^{d_i}.$$

We then set $u := u_M$ and $P := G_M \prod_{i=1}^{N}(T - \alpha_i)^{d_i}$ to conclude. □

Lemma 3.12. *Let X be a quasi-smooth k-analytic space of dimension 2.*

1. *Let $q \in X_{rig}$ and $\pi : \tilde{X} \to X$ the blowing-up of X at q, and let $S \subseteq \tilde{X}$ be a locally semianalytic subset. Then $\pi(S)$ is locally semianalytic.*
2. *If $\pi : \tilde{X} \to X$ is a succession of blowing-up of rigid points, and $S \subseteq \tilde{X}$ is locally semianalytic, then $\pi(S)$ is also locally semianalytic.*

Proof. (2) is a consequence of (1) so we only have to show (1).
The problem is local on X, and since outside q, π is a local isomorphism, we can restrict to an affinoid neighbourhood of q, and since X is regular at q, we can assume that $X = \mathbb{B}^2$ and q is the origin.
Then $\pi : \tilde{X} \to X$ can be described with two charts, one of them being

$$\begin{array}{rrcl} \pi_1 : & X_1 = \mathcal{M}(k\{x, t\}) & \to & X = \mathcal{M}(k\{x, y\}) \\ & (x, t) & \mapsto & (x, tx) \end{array}$$

The other chart being analogous we only consider π_1. Now, changing S in $S \cap X_1$, we can restrict to show that if S is locally semianalytic in X_1, so is $\pi_1(S)$. Since π_1 induces an isomorphism between $X_1 \setminus V(x)$ and $\{p \in \mathbb{B}^2 \mid |y(p)| \leq |x(p)| \neq 0\}$, we only have to show that $\pi_1(S)$ is semianalytic around the origin q.
Now if for each $p \in \mathbf{E} := V(x) \subseteq X_1$ we can find V_p an affinoid neighbourhood of p, and $\varepsilon_p > 0$ such that $\pi_1(V_p \cap S) \cap \mathbb{B}^2_{\varepsilon_p}$ is globally semianalytic in $\mathbb{B}^2_{\varepsilon_p} \subseteq X$, then by compactness, we can extract $V_1, \ldots, V_n$ a finite covering of $\mathbf{E}$ and $\varepsilon > 0$ such that $\cup_{i=1}^{n} (\pi_1(V_i \cap S)) \cap \mathbb{B}^2_\varepsilon = \pi_1(S) \cap \mathbb{B}^2_\varepsilon$ is globally semianalytic. So we fix $p \in \mathbf{E} = V(x)$ and try to find V an affinoid neighbourhood of p, and $\varepsilon > 0$ such that $\pi_1(V \cap S) \cap \mathbb{B}^2_\varepsilon$ is globally semianalytic in $\mathbb{B}^2_\varepsilon$.
Since S is locally semianalytic in X_1, we can find V an affinoid neighbourhood of p such that $V \cap S$ is globally semianalytic in V. According to corollary 3.2, we can assume that[5] $V = \mathbb{B}_\varepsilon \times W$ where

$$W = \{w \in \mathcal{M}(k\{t\}) \mid |(t - a_0)(w)| \leq r_0 \text{ and } |(t - a_i)(w)| \geq r_i,\ i = 1 \ldots n\}$$

with $a_0, \ldots, a_n \in k$.
To simplify the notations, we can also assume that the globally semianalytic subset S of V we are dealing with is of the following form:

$$S = \bigcap_{j=1}^{m} \{v \in V \mid |f_j(v)| \Diamond_j |g_j(v)|\}$$

Now remind that $V = \mathbb{B}_\varepsilon \times W$ with $\mathbb{B}_\varepsilon = \mathcal{M}(k\{\varepsilon^{-1}x\})$. So we can factor each f_j and g_j by the greatest power of x which is a factor, hence introduce some integers b_j, c_j such that

$$S = \bigcap_{j=1}^{m} \{v \in V \mid |x^{b_j} \tilde{f}_j(v)| \Diamond_j |x^{c_j} \tilde{g}_j(v)|\}$$

where the series $\tilde{f}_j(0, t)$ and $\tilde{g}_j(0, t)$ are non zero, and $f_j = x^{b_j} \tilde{f}_j, g_j = x^{c_j} \tilde{g}_j$. But to simplify the notations, we'll use f_j (resp. g_j) instead of $\tilde{f}_j$ (resp. $\tilde{g}_j$), so that

$$S = \bigcap_{j=1}^{m} \{v \in V \mid |x^{b_j} f_j(v)| \Diamond_j |x^{c_j} g_j(v)|\}$$

where the series $f_j(0, t)$ and $g_j(0, t)$ are non zero.
Then according to lemma 3.10 we can decrease ε and W so that for each $f_j, g_j \in \{f_1 \ldots f_m, g_1, \ldots, g_m\}$, $f_j = u_j P_j$ (resp. $g_j = v_j Q_j$) where u_j (resp. v_j) is a multiplicative unit, and P_j (resp. Q_j) $\in k\{\varepsilon^{-1}x\}[t, (t - a_1)^{-1}, \ldots, (t - a_n)^{-1}]$.

---

[5] Here we use the explicit description of affinoid domains of $\mathbb{B}$.

Said differently, and with different notations, there exists an integer N such that $f_j = u_j \cdot \frac{P_j}{((t-a_1)\dots(t-a_n))^N}$ where u_j is a multiplicative unit and $P_j \in k\{\varepsilon^{-1}x\}[t]$ (and resp. $g_j = v_j \cdot \frac{Q_j}{((t-a_1)\dots(t-a_n))^N}$). Hence

$$|f_j(v)|\Diamond_j|g_j(v)| \Leftrightarrow \left| u_j(v)\frac{P_j(v)}{((t-a_1)\dots(t-a_n))^N(v)}\right| \Diamond_j \left| v_j(v)\frac{Q_j(v)}{((t-a_1)\dots(t-a_n))^N(v)}\right| \Leftrightarrow |P_j(v)|\Diamond_j\lambda_j|Q_j(v)|$$

where

$$\lambda_j = \frac{\|v_j\|}{\|u_j\|} \in |k^\times|.$$

Moreover,

$$S\cap V = (S\cap\{v\in V \mid x(v)=0\})\cup(S\cap\{v\in V \mid x(v)\neq 0\})$$

and $\pi_1(\{v\in V \mid x(v)=0\}) = q$, the origin of $\mathbb{B}^2$.

So, adding if necessary the origin to $\pi_1(S\cap\{v\in V \mid v(x)\neq 0\})$ (which won't change the fact that it is globally semianalytic), we can restrict to show that $\pi_1(S\cap\{v\in V \mid v(x)\neq 0\})$ is globally semianalytic around the origin. Moreover since on $\{v\in V \mid v(x)\neq 0\}$, π_1 is bijective, the following holds:

$$\pi_1\left(\bigcap_{j=1}^m \{v\in V \mid |x^{b_j}f_j(v)|\Diamond_j|x^{c_j}g_j(v)|\}\cap\{v\in V \mid x(v)\neq 0\}\right)$$

$$=\bigcap_{j=1}^m \pi_1\left(\{v\in V \mid |x^{b_j}f_j(v)|\Diamond_j|x^{c_j}g_j(v)|\}\right)\cap\{v\in V \mid x(v)\neq 0\}$$

Now since $y=tx$ and $P_j\in k\{\varepsilon^{-1}x\}[t]$ there exists an integer $M\geq 0$ such that $x^MP_j(x,t)\in k\{\varepsilon^{-1}x\}[tx] = k\{\varepsilon^{-1}x\}[y]$, i.e. $x^MP_j(x,t) = \pi^*(\tilde{P}_j(x,y))$ for some $\tilde{P}_j(x,y)\in k\{\varepsilon^{-1}x\}[y]$ and such that $x^MQ_j(x,t)\in k\{\varepsilon^{-1}x\}[y]$, i.e. $x^MQ_j(x,t)=\pi^*(\tilde{Q}_j(x,y))$ for some $\tilde{Q}_j(x,y)\in k\{\varepsilon^{-1}x\}[y]$.
Now on $\{v\in V \mid v(x)\neq 0\}$,

$$|x^{b_j}f_j(v)|\Diamond_j|x^{c_j}g_j(v)| \Leftrightarrow |x^{M+b_j}f_j(v)|\Diamond_j|x^{M+c_j}g_j(v)| \Leftrightarrow |x^{b_j}\tilde{P}_j(\pi_1(v))|\Diamond_j\lambda_j|x^{c_j}\tilde{Q}_j(\pi_1(v))|.$$

From that we conclude that

$$z\in\pi_1\left(\bigcap_{j=1}^m \{v\in V \mid |x^{b_j}f_j(v)|\Diamond_j|x^{c_j}g_j(v)|\}\cap\{v\in V \mid x(v)\neq 0\}\right)$$

$$\Leftrightarrow z\in\bigcap_{j=1}^m\{z\in\pi_1(V) \mid |x^{b_j}\tilde{P}_j(z)|\Diamond_j|x^{c_j}\tilde{Q}_j(z)|\}\cap\{z\in X \mid x(z)\neq 0\}$$

Since $\pi_1(\mathbb{B}^2_\varepsilon)\subseteq\mathbb{B}^2_\varepsilon$ and is globally semianalytic in $\mathbb{B}^2_\varepsilon$, we conclude that $\pi_1(S\cap\{v\in V \mid v(x)\neq 0\})$ is globally semianalytic in $\mathbb{B}^2_\varepsilon$, which ends the proof. □

Theorem 3.13. *Let X be a quasi-smooth k-analytic space of dimension 2 with k algebraically closed, and $S\subseteq X$. Then S is overconvergent subanalytic subset if and only if S is locally semianalytic.*

Proof. Since the problem is local, we can assume that X is affinoid and that $S=\varphi(U)$ where $(Y,U)\overset{\varphi}{\dashrightarrow}X$ is a constructible datum, and just check that S is locally semianalytic. We do it by induction on the complexity of φ. So let $(Y,U)\overset{\varphi}{\dashrightarrow}X$ be a constructible datum, that we decompose as

$$\varphi = (Y,U)\overset{\psi}{\dashrightarrow}Z\xrightarrow{\chi}X$$

where χ is an elementary constructible datum, and ψ a constructible datum whose complexity is one less than φ. So we can introduce $f,g\in\mathcal{A}$, $0<s<r$ such that $Z=\mathcal{M}(\mathcal{A}\{r^{-1}t\}/(f-tg))$. According to proposition 3.9, we can find a succession of blowing-up of rigid points $\pi:\tilde{X}\to X$

such that for all $x \in \tilde{X}$, $f_x | g_x$ or $g_x | f_x$. According to remark 3.8, $\tilde{X}$ is still quasi-smooth. This gives us the following cartesian diagram:

$$\begin{tikzcd}
(Y,U) \arrow[r, dotted, "\varphi"] & X \\
(Y',U') \arrow[u, "\pi'"] \arrow[r, dotted, "\varphi'"] & \tilde{X} \arrow[u, "\pi"']
\end{tikzcd}$$

Then $\varphi(U) = \pi(\varphi'(U'))$. Moreover, since $\tilde{X}$ is compact, we can then find a finite wide covering $\{X_i\}_{i=1}^n$ of $\tilde{X}$ by affinoid domains such that on each X_i, $f|g$ or $g|f$, here we see f and g as functions on X_i. We denote by $\pi_i : X_i \to X$ the composition of the embedding of the affinoid domain $X_i \to \tilde{X}$ with $\pi : \tilde{X} \to X$. This gives the following cartesian diagrams:

$$\begin{tikzcd}
(Y,U) \arrow[r, dotted, "\psi"] & Z \arrow[r, "\chi"] & X \\
(Y_i,U_i) \arrow[u, "\pi_i''"] \arrow[r, dotted, "\psi_i"] & Z_i \arrow[u, "\pi_i'"] \arrow[r, "\chi_i"] & X_i \arrow[u, "\pi_i"]
\end{tikzcd}$$

Then

$$\varphi(U) = \pi(\varphi'(U')) = \pi\left(\bigcup_{i=1}^n \chi_i(\psi_i(U_i))\right)$$

But $(Y_i, U_i) \overset{\psi_i}{\dashrightarrow} Z_i$ is a constructible datum of lower complexity than φ, so that we would like to use our induction hypothesis, and claim that $\psi_i(U_i)$ is locally semianalytic. However, Z_i is not necessarily still quasi-smooth so we can't do that. However, according to remark 3.7, we can in fact replace Z_i and in fact assume that Z_i is a Weierstrass (or a Laurent) domain of X_i, and hence is quasi-smooth. Hence by induction hypothesis $\psi_i(U_i)$ is locally semianalytic in Z_i. Now we use lemma 3.6 to assert that $\chi_i(\psi_i(U_i))$ is locally semianalytic in X_i. So

$$\varphi'(U') = \bigcup_{i=1}^n (\chi_i(\psi_i(U_i))$$

is locally semianalytic in $\tilde{X}$, since $\{X_i\}$ was a wide covering of $\tilde{X}$. Finally, according to lemma 3.12, $\pi(\varphi'(U')) = S$ is also locally semianalytic. $\square$

References

[1] V.G. Berkovich. *Spectral theory and analytic geometry over non-Archimedean fields.* Amer Mathematical Society, 1990.

[2] V.G. Berkovich. Etale cohomology for non-Archimedean analytic spaces. *Publications Mathématiques de l'IHÉS*, 78(1):5–161, 1993.

[3] S. Bosch, U. Güntzer, and R. Remmert. *Non-Archimedean analysis*, volume 261 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences].* Springer-Verlag, Berlin, 1984. A systematic approach to rigid analytic geometry.

[4] S. Bosch and W. Lutkebohmert. Formal and rigid geometry. I : Rigid Spaces. *Mathematische Annalen*, 295(3):291–317, 1993.

[5] R. Cluckers and L. Lipshitz. Fields with analytic structure. *J. Eur. Math. Soc. (JEMS)*, 13(4):1147–1223, 2011.

[6] J. Denef and L. van den Dries. p-adic and real subanalytic sets. *Ann. of Math. (2)*, 128(1):79–138, 1988.

[7] A. Ducros. Flatness in non-Archimedean analytic geometry. *preprint, arXiv:[1107.4259].*

[8] Antoine Ducros. Parties semi-algébriques d'une variété algébrique p-adique. *Manuscripta Math.*, 111(4):513–528, 2003.

[9] Jean Fresnel and Marius van der Put. *Rigid analytic geometry and its applications*, volume 218 of *Progress in Mathematics.* Birkhäuser Boston Inc., Boston, MA, 2004.

[10] T. S. Gardener and Hans Schoutens. Flattening and subanalytic sets in rigid analytic geometry. *Proc. London Math. Soc. (3)*, 83(3):681–707, 2001.

[11] T. S. Gardener and Hans Schoutens. Rigid subanalytic sets. In *Valuation theory and its applications, Vol. I (Saskatoon, SK, 1999)*, volume 32 of *Fields Inst. Commun.*, pages 141–150. Amer. Math. Soc., Providence, RI, 2002.

[12] R. Hartshorne. *Algebraic geometry.* Springer, 2000.

[13] Roland Huber. *Étale cohomology of rigid analytic varieties and adic spaces.* Aspects of Mathematics, E30. Friedr. Vieweg & Sohn, Braunschweig, 1996.

[14] János Kollár. *Lectures on resolution of singularities*, volume 166 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2007.

[15] Serge Lang. *Algebra*, volume 211 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, third edition, 2002.

[16] L. Lipshitz. Rigid subanalytic sets. *Amer. J. Math.*, 115(1):77–108, 1993.

[17] L. Lipshitz and Z. Robinson. Flattening and analytic continuation of affinoid morphisms. Remarks on a paper of T. S. Gardener and H. Schoutens: "Flattening and subanalytic sets in rigid analytic geometry" [Proc. London Math. Soc. (3) **83** (2001), no. 3, 681–707; mr1851087]. *Proc. London Math. Soc. (3)*, 91(2):443–458, 2005.

[18] Leonard Lipshitz and Zachary Robinson. Rings of separated power series and quasi-affinoid geometry. *Astérisque*, (264):vi+171, 2000.

[19] Hans Schoutens. Rigid subanalytic sets. *Compositio Math.*, 94(3):269–295, 1994.

[20] Hans Schoutens. Rigid subanalytic sets in the plane. *J. Algebra*, 170(1):266–276, 1994.

[21] Hans Schoutens. Uniformization of rigid subanalytic sets. *Compositio Math.*, 94(3):227–245, 1994.

[22] John Tate. Rigid analytic spaces. *Invent. Math.*, 12:257–289, 1971.

Institut de Mathématiques de Jussieu, Université Pierre-et-Marie-Curie, 4 Place de Jussieu, 75005 Paris

E-mail address: `fmartin@math.jussieu.fr`